

Clinical Profile of Neonates with Hypernatremic Dehydration Hospitalized in an Outborn Neonatal Intensive Care Unit

SMITA NAIR, AMITABH SINGH AND *MAMTA JAJOO

Department of Pediatrics, Chacha Nehru Bal Chikitsalaya, New Delhi, India.

**mamtajajoo123@gmail.com*

PII: S097475591600111

Note: This early-online version of the article is an unedited manuscript that has been accepted for publication. It has been posted to the website for making it available to readers, ahead of its publication in print. This version will undergo copy-editing, typesetting, and proofreading, before final publication; and the text may undergo minor changes in the final version.

ABSTRACT

This retrospective observational study aimed to find the clinical profile of hypernatremic dehydration in neonates. 49 neonates (3.4% of the total admitted newborns) developed hypernatremic dehydration. The major presenting complaints were fever (34.6%), poor feeding (42.8%), loose stools (40.8%) and lethargy (26.5%). The time needed for correction of hypernatremia was 38.6 (\pm 15.1) hours. Exclusively breastfed neonates had lesser complication rates of hypernatremic dehydration.

Keywords: *Breastfeeding, Dehydration, Neonatal, Sepsis.*

Hypernatremic dehydration is a common but life threatening condition in neonates. Inadequate breastfeeding, improperly diluted mixed feeding, and gastrointestinal losses are the main etiologies for hypernatremia in neonates. Hypernatremia may cause intracerebral edema, hemorrhage, and gangrene, resulting in death or long-term morbidity [1,2]. Data on management and outcome of hypernatremic dehydration from developing world is limited. There is no universally accepted fluid regimen for management of hypernatremic dehydration [3].

Between January 2014 to August 2015, 1510 neonates were admitted in the out born neonatal intensive care unit of a pediatric hospital situated in Northern India. Chart review of neonates with moderate and severe hypernatremic dehydration (serum sodium >150 meq/L) was done. Neonates with documented hypernatremia and presenting without any prior treatment were included; those with proven sepsis were excluded. Hypernatremia was corrected using formula based on water deficit and solute deficit [4].

Forty-nine neonates were admitted with hypernatremic dehydration. The presenting complaints were fever (34.6%), poor feeding (42.8%), loose stools (40.8%) and lethargy (26.5%), decreased urine output (8.2%) weight loss (75.5%). 24.5% neonates presented with neurological complaints and examination revealed a doughy feel of skin in 90 % of the neonates. Thirty-three (67.3%) neonates were hospital delivered and 6 (12.2%) had history of birth asphyxia. Seven neonates (14.8%) required ionotropes and five had culture positive sepsis. The time needed for correction of hypernatremia was 38.6 (\pm 15.1) hours. Mean percentage of dehydration on presentation was 16.3 (SD 11.03). Mean sodium on admission was 157.7 (SD 9.41) mEq/L. Hyperkalemia and metabolic acidosis was present in 21 (42.8%) and 39 (79.6%) neonates, respectively. The duration of hospital stay was 7.1 (\pm 4.8) days. Clinical and laboratory characteristics of neonates with hypernatremia compared to feeding status is shown in **Table I**.

The incidence of hypernatremic dehydration in present study was 3.4% with a female preponderance. Poor feeding was the most common presenting complaint. Exclusively top fed neonates

had higher percentage of acute kidney injury, mean sodium level, mean creatinine value at presentation and were more dehydrated compared to other groups.

Oddie, *et al.* [4] have reported an incidence of hypernatremic dehydration as 2.5/10000 live births and Moritz, *et al.* [5] found a 5-year incidence of breastfeeding associated hypernatremia among hospitalized neonate to be 1.9%. The mean age of presentation was 14.8 (± 8.3) days which is comparable to previous studies (4-21 days). We found 61.2% neonates with $>10\%$ weight loss which is comparable to study by Uras, *et al.* [5].

We did not find caesarian section (18.3%) to be a risk factor for hypernatremic dehydration as has been reported in previous studies [6,7]. A large number of home deliveries in the study population may account for this difference. Our study confirmed more cases of hypernatremic dehydration in primigravida mothers (46.9%) as reported by Moritz, *et al.* [8].

The time taken to correct hypernatremia was 38.6 (± 15.1) hours. Faith, *et al.* found a fall in sodium levels of 0.48 (± 0.2) mEq/L in first 24 hours and 0.4 (± 0.3) mEq/L in the next 24 hours [9]. At the time of discharge neurological examination was normal in 85.7% neonates. Limitations of present study include its retrospective design, lack of neurodevelopment follow up, lack of correlation between breast milk sodium levels and hypernatremia, and that the correction rate in serum sodium levels in first 6 and 24 hours was not recorded.

Poor feeding is both a cause and a manifestation of hypernatremic dehydration. Exclusively breastfed neonates with hypernatremic dehydration have fewer complications. Fluid management is cornerstone of good outcome in hypernatremic dehydration and such cases can be managed at level II neonatal intensive care unit [10].

Contributors: MJ: Concept and designed the study, analyzed data and drafted the manuscript; SN: Collected the data and helped in data analysis, drafted the manuscript; AS: collected and analyzed the data, reviewed the literature, drafted the manuscript.

Funding: None; *Competing interest:* None stated.

REFERENCES

1. Moritz ML, Ayus JC. Preventing neurological complications from dysnatremias in children. *Pediatr Nephrol.* 2005;20:1687-700.
2. Moritz ML, Ayus JC. The changing pattern of hypernatremia in hospitalized children. *Pediatrics.* 1999;104:435-9.
3. Maayan-Metzger A, Mazkereth R, Kuint J. Fever in healthy asymptomatic newborns during the first days of life. *Arch Dis Child Fetal Neonatal Ed.* 2003;88:F312-4.
4. Oddie S, Richmond S, Coulthard M. Hypernatraemic dehydration and breast feeding: A population study. *Arch Dis Child.* 2001;85:318-20.
5. Uras N, Karadag A, Dogan G, Tonbul A, Tatli MM. Moderate hypernatremic dehydration in newborn infants: retrospective evaluation of 64 cases. *J Matern Fetal Neonatal Med.* 2007;20:449-52.
6. Manganaro R, Mami C, Marrone T, Marseglia L, Gemelli M. Incidence of dehydration and hypernatremia in exclusively breast-fed infants. *J Pediatr.* 2001;139:673-5.
7. Erdemir A, Kahramaner Z, Cosar H, Turkoglu E, Kanik A, Sutcuoglu S, *et al.* Comparison of oral and intravenous fluid therapy in newborns with hypernatremic dehydration. *J Matern Fetal Neonatal Med.* 2014;27:491-4.
8. Moritz ML, Manole MD, Bogen DL, Ayus JC. Breastfeeding-associated hypernatremia: are we missing the diagnosis? *Pediatrics.* 2005;116:e343-7.
9. Bolat F, Oflaz MB, Güven AS, Özdemir G, Alaygut D, Doğan MT, *et al.* What is the safe approach for neonatal hypernatremic dehydration? A retrospective study from a neonatal intensive care unit. *Pediatr Emerg Care.* 2013;29:808-13.
10. Alshayeb HM, Showkat A, Babar F, Mangold T, Wall BM. Severe hypernatremia correction rate and mortality in hospitalized patients. *Am J Med Sci.* 2011;341:356-60.

TABLE I CLINICAL AND LABORATORY CHARACTERISTICS IN NEONATES WITH HYPERNATREMIA CLASSIFIED AS PER BREAST FEEDING STATUS

<i>Parameter</i>	<i>Exclusively breastfed (n=16)</i>	<i>Exclusively top fed (n=17)</i>	<i>Mixed fed (n=16)</i>
Acute Kidney Injury	11 (68.7%)	14 (82.3%)	8 (50%)
Serum Sodium (meq/L) Mean (SD) Median (IQR)	150.1 (9.7) 150 (148-159)	162.3 (11) 164.5 (145-165)	155 (10.7) 158 (149-163)
Serum Potassium (meq/L) Mean (SD) Median(IQR)	6 (1.3) 5.3 (4-6)	5 (1.3) 4.9 (4-5.9)	10 (1.3) 6.9 (5-7.1)
Serum Creatinine Mean (SD) Median (IQR)	2.2 (0.98) 2.5 (2-4.2)	3.5 (0.9) 4.1 (2-5.1)	2.7 (0.8) 3.1 (2.5-5.2)
Correction time for Hypernatremia(h) Mean (SD) Median(IQR)	33.6 (15.1) 30 (20-40)	44.6 (17.2) 42 (24-56)	37.7 (17) 30 (24-42)
Duration of hospitalization (d) Mean (SD) Median(IQR)	6.3(4.8) 6 (4.75-8)	8.8 (4.6) 7 (4.5-8.75)	6.1 (4.5) 6.5 (4.1-8)
Neonates with signs of dehydration	7.5%	9.2%	7.9%