

WEB TABLE I SUBGROUP - ANALYSIS OF INFANTS WITH AND WITHOUT BREAKTHROUGH SEIZURES IN EITHER GROUP

<i>No breakthrough seizures (n=98)</i>	<i>Breakthrough seizures (n=54)</i>	<i>P value</i>
<i>HIE (at admission)*</i>		
Stage II	63/92 (68.47)	38/50 (76.00)
Stage III	13/92 (14.13)	10/50 (20.00)
Serum phenobarbitone level (mcg/mL) [#] at 12 h, mean (SD)	22.15 (21.87)	22.91 (22.65)
Infants with phenobarbitone level* <10 mcg/mL	28 (28.5)	16 (29.3)
Time to full enteral feeds [#] (d)	5.25 (3.44)	7.07 (3.43)
Duration of hospital stay [#] (d)	6.46 (3.67)	8.87 (4.77)
<i>Abnormal Neurological examination*</i>		
At discharge	20/98 (20.4)	19/54 (35.18)
At 1 month	9/82 (10.98)	12/44 (27.27)
At 3 month	1/73 (1.37)	3/35 (8.57)
<i>Seizure recurrence*</i>		
1 month	5/82 (6.1)	4/44 (9.1)
3 month	5/73 (6.8)	4/35 (11.4)
Re-hospitalization rate*	9/82 (10.91)	8/44 (18.18)

HIE: Hypoxic ischemic encephalopathy. Value indicate mean (SD)[#], or n (%)*.