

**WEB TABLE I** CLINICAL CHARACTERISTICS OF SRNS CHILDREN WITH AND WITHOUT SUBCLINICAL OR OVERT HYPOTHYROIDISM

<i>Variable</i>	<i>SRNS with subclinical or overt hypothyroidism (n=10)</i>	<i>SRNS without subclinical or overt hypothyroidism (n=20)</i>	<i>P value</i>
Age of onset of SRNS (y)	5.8 (3.3)	8 (4.0)	0.172
Age of onset of NS (y)	3.2 (3.0)	5.8 (3.6)	0.066
Initial resistance*	3 (30%)	9 (45%)	0.694
Duration of onset of SRNS to thyroid status evaluation (mo)	10 (14.6)	9.9 (12.0)	0.976
Total duration of illness (y)	2.7 (2.6)	2.1 (1.3)	0.366
Serum albumin (g/dL)	2.2 (0.8)	2.8 (1.0)	0.124
Serum cholesterol (mg/dL)	392.8 (128.1)	364.5 (213.6)	0.700
<i>Histopathology*</i>			
FSGS	2 (20%)	11 (55%)	0.373
MCD	3 (30%)	6 (30%)	
MesPGN	5 (50%)	3 (15%)	

*Values are expressed in mean (SD) or \*n(%); FSGS-Focal segmental glomerulosclerosis; MCD-Minimal Change Disease; MesPGN-mesangioproliferative glomerulonephritis; SRNS-steroid resistant nephrotic syndrome.*