

## Transport-related Adverse Events in Critically-ill Children: The Role of a Dedicated Transport Team

SUMANT PRABHUDESAI, MOHANBABU KASALA, NITIN MANWANI, RAVIKUMAR KRUPANANDAN AND BALA RAMACHANDRAN

From Department of Paediatric Critical Care, Kanchi Kamakoti CHILDS Trust Hospital, Chennai, India.

### Correspondence to:

Dr Sumant Prabhudesai,

Department of Paediatric Critical Care, Kanchi Kamakoti CHILDS Trust Hospital, 12A, Nageswara Road, Nungambakkam, Chennai, India.  
sumantprabhudesai2014@gmail.com

Received: July 22, 2016;

Initial Review: December 13, 2016;

Accepted: July 22, 2017.

**Objective:** To compare the frequency of transport-related adverse events in children during specialized, non-specialized or unassisted transports. **Methods:** Patients were grouped based on transport team involved – specialized (Group-1); non-specialized (Group-2); unassisted transport (Group-3). Demographics, events during transport and condition on arrival were recorded. **Results:** Group-1 children had a lower incidence of adverse events compared to Group-2 and Group-3 (4.3%, 82.6% and 85.4% respectively;  $P < 0.001$ ). At arrival, children in Group-1 had a lower incidence of respiratory distress and airway compromise ( $P < 0.001$ ). **Conclusion:** Transport of critically ill children by a specialized transport team is associated with fewer transport-related adverse events.

**Keywords:** Airway compromise, Emergency care, Intubation, Respiratory failure.

Published online: August 24, 2017. PII:S097475591600081

Medical transport is a key element in the chain of survival for critically ill children, especially in resource-limited settings where critical care services are scarce. The challenges in pediatric transports differ considerably from those in adults. Limited training with pediatric patients may restrict the ability of general transport teams to transfer seriously ill children. Evidence shows that children transported by specialized teams have fewer Transport-related adverse events (TRAEs), a more stable physiology upon arrival, and lower in-hospital mortality [1-5]. In India, specialized pediatric transport teams are few [6], and data on transport practices are scarce [7]. We conducted this study to determine whether inter-hospital transport by specialized transport teams reduces the incidence of TRAEs in children in comparison to non-specialized teams and unassisted transports.

### METHODS

This prospective observational study was conducted at a tertiary care children's referral hospital in Chennai, India from February 2014 to September 2014. Approval was obtained from the Hospital Ethics Committee.

All patients aged between 1 month and 18 years who were transported to the hospital's Emergency Room (ER) and required Pediatric intensive care unit (PICU) admission were included. Depending on the mode of

transportation, they were assigned to one of three groups: Group-1: transported by a specialized pediatric transport team; Group-2: transported by a general transport team; or Group-3: brought by caregivers without medical assistance. A specialized team was defined as one which: (i) was attached to a hospital having a dedicated children's transport team; (ii) had a doctor trained or undergoing training in pediatrics, pediatric anesthesia or pediatric critical care, and skilled in airway management and cardiopulmonary resuscitation; (iii) had a nurse with pediatric experience; and (iv) had an ambulance equipped for emergency airway management, vascular access, oxygen, drug delivery and a multi-parameter monitor. Transport teams not meeting all these criteria were considered general transport teams.

Data on demographics, transport-related adverse events, interventions during transport, condition and interventions at arrival to ER were collected. The primary outcome measured was the occurrence of TRAEs. A TRAE included hypoxia, airway compromise requiring intervention, pneumothorax, tachycardia, bradycardia, hypotension, cardiac arrest or hypoglycemia (all defined as per PALS guidelines [8]), which were recognized during transport or immediately after arrival in ER. The secondary outcomes measured were PICU length of stay and mortality. Severity of illness was assessed by the Pediatric Risk of Mortality (PRISM) III score.

*Statistical analysis:* Student's t test was used for parametric data and Kruskal-Wallis test for non-parametric data. Fisher's exact test was used for categorical variables. Variables significant on univariate analysis were further analyzed by logistic regression. A two-tailed *P* value of <0.05 was considered significant.

## RESULTS

We included, 204 children; majority (58.3%) in Group-3. In Group-2, 9 (23.1%) patients were accompanied by a doctor while 30 (76.9%) were accompanied by paramedical staff. In Group-3, 15 (12.6%) children were brought by ambulance without paramedical staff while the other 104 (87.4%) were brought by private vehicles.

Children who were younger ( $P=0.003$ ), had a respiratory problem ( $P=0.03$ ), or required transportation over distances less than 5 km ( $P=0.01$ ) were more often transported by specialized teams than non-specialized ones (**Table I**).

TRAEs occurred in 142 (69.6%) children (**Table II**). Group-1 children had a significantly lower incidence of TRAEs compared to Group-2 and Group-3 (4.3%, 82.6%

and 85.4% respectively,  $P<0.001$ ). Airway compromise, hypoxia and tachycardia were less common in Group-1 compared to Group-2 and Group-3 ( $P<0.001$ ).

Group-1 children had a lower incidence of respiratory distress ( $P<0.001$ ), requirement of oxygen support ( $P<0.001$ ) and emergency intubation ( $P<0.001$ ) upon arrival to ER. The length of stay and mortality were not significantly different between the three groups.

On univariate analysis, neurological problems, transport distance >5 km, non-specialized/unassisted transport and transport time >3 hours were identified as risk factors for TRAEs. On multivariate logistic regression, transport time >3 hours ( $P=0.002$ ) and non-specialized/unassisted transport ( $P<0.001$ ) were independent risk factors for TRAEs. The relative risk of TRAEs with non-specialised/unassisted transport was 20.4.

## DISCUSSION

In this observational study, we observed that the large majority of pediatric transports occurred unassisted or by non-specialized teams wherein the risks of airway compromise, respiratory distress and tachycardia were high.

**TABLE I** CHARACTERISTICS OF CHILDREN TRANSPORTED TO THE EMERGENCY DEPARTMENT ( $N=204$ )

	Total	Group-1(n= 46)	Group-2 (n= 39)	Group-3(n= 119)	<i>P</i> value
Age* (mo)	19.5 (9- 69.2)	12 (5.2- 25.8)	30 (10-92)	24 (9- 72)	0.003
Females	80 (39.2)	18 (39)	14 (35.9)	48 (40.3)	0.88
PRSIM III score*	3 (0.75- 6)	3 (0- 7)	3 (2- 6)	3 (1.5- 5.5)	0.98
<i>Indication for transport</i>					
Neurological	76 (37.3)	14 (30.4)	20 (51.3)	42 (35.3)	0.03
Respiratory	85 (41.7)	23 (50)	14 (35.9)	48 (40.3)	
Metabolic	18 (8.8)	0 (0)	1 (2.6)	17 (14.3)	
Shock	12 (5.9)	4 (8.7)	1 (2.6)	7 (5.9)	
Trauma	13 (6.4)	5 (10.9)	3 (7.6)	5 (4.2)	
<i>Distance (km)</i>					
<5	28 (13.7)	12 (26.1)	2 (5.1)	14 (11.8)	0.01
5-10	59 (28.9)	8 (17.4)	8 (20.5)	43 (36.1)	
10-20	30 (14.7)	7 (15.2)	6 (15.4)	17 (14.3)	
>20	87 (42.6)	19 (41.3)	23 (58.9)	45 (37.8)	
<i>Transport support</i>					
Private vehicle	105 (51.5)	0 (0)	0 (0)	104 (87.4)	
Ambulance	99 (48.5)	46 (100)	39 (100)	15 (12.6)	
Doctor	55 (26.9)	46 (100)	9 (23.1)	0 (0)	
Paramedic	76 (37.3)	46 (100)	30 (76.9)	0 (0)	

Group-1: transport by specialised teams; Group-2: transport by general/unspecialised teams teams; Group-3: unassisted transport. Variables are expressed as \*median (IQR) or numbers (percentage); PRISM: Pediatric Risk of Mortality.

**TABLE II** TRANSPORT-RELATED ADVERSE EVENT (TRAE), EMERGENCY INTERVENTIONS AND OUTCOME IN STUDY CHILDREN (N=204)

TRAEs	Total	Group-1(n= 46)	Group-2(n= 39)	Group-3(n= 119)	P value
At least 1 TRAE	142 (69.6)	2 (4.3)	38 (97.4)	102 (85.7)	<0.001
Hypoxia	81 (39.7)	0 (0)	24 (61.5)	57 (47.9)	<0.001
Hypotension	14 (6.9)	1 (2.2)	3 (7.7)	10 (8.4)	0.36
Tachycardia	100 (49)	0 (0)	26 (66.6)	74 (62.2)	<0.001
Hypoglycemia	3 (1.5)	1 (2.2)	1 (2.6)	1 (0.84)	0.67
Airway compromise	53 (25.9)	0 (0)	19 (48.7)	34 (28.6)	<0.001
<i>Condition at arrival</i>					
Shock	47 (23)	11 (23.9)	6 (15.4)	30 (25.2)	0.44
Respiratory distress	79 (38.7)	4 (8.7)	23 (58.9)	52 (43.7)	<0.001
Low GCS(<8/15)	40 (19.6)	6 (13)	10 (25.6)	24 (20.2)	0.34
<i>Emergency interventions at arrival</i>					
Oxygen	92 (45.1)	5 (10.9)	16 (41)	71 (59.7)	<0.001
Intubation/ re-intubation	49 (24)	1 (2.2)	17 (43.6)	31 (26.1)	<0.001
Fluid bolus	68 (33.3)	12 (26.1)	9 (23.1)	47 (39.5)	0.09
Inotropes	20 (9.8)	6 (13)	4 (10.3)	10 (8.4)	0.58
<i>Outcome</i>					
*PICU length of stay (d)	4 (3- 6)	4 (2- 7)	5 (3- 7)	4 (3- 5)	0.24
Death	35 (17.1)	8 (17.4)	9 (23.1)	18 (15.1)	0.52

Transport by: Group-1: specialized teams; Group-2: general/unspecialized teams; Group-3: unassisted transport. Variables are expressed as n(%) or \*median (IQR); GCS: Glasgow Coma Scale.

Having no prior data on existing transport practices, we could not calculate a sample size; therefore the study duration and patient recruitment was arbitrary. The unavailability of pre-transport clinical details was a major limitation. The large difference in the number of patients per group resulting from recruitment of consecutive patients presenting to the ER may well have affected statistical analysis. This difference largely reflects the prevalent referral practices in the region.

Systematic information on pediatric transport practices in India is scarce. A recent study from Delhi reported that the majority of children referred to the ER from another facility arrived by ambulance unassisted [7]. A retrospective analysis of records in neonates showed nearly 45% of the babies were transported by paramedical or non-medical persons [9]. In another study, only 29% of neonates were transported by ambulance, the others being brought by private vehicles or public transport [10].

Worldwide, data on pediatric transport are mainly available from countries with established emergency transport systems. North American studies have shown that adverse events were less when children were accompanied by a tertiary-care physician and higher

when team members had no pediatric transport training [2,3]. Transportation by non-specialized teams was associated with more unplanned events and higher mortality [5,11,12]. We did not find any difference in mortality in our study.

Kumar, *et al.* [13] documented that distance or duration of transport did not affect the risk of adverse events and outcomes in neonates transported by a qualified transport team [13]. In our study, the greater risk associated with longer durations likely reflects the fact that most long distance transports were unassisted or done by non-specialized teams.

In conclusion, utilizing specialized pediatric transport teams can reduce the incidence of adverse effects during transport. Further large-scale and multicentric studies are necessary to determine whether this translates to reduced morbidity and mortality.

*Acknowledgement:* The CHILDS Trust Medical Research Foundation for their support in conducting this study.

*Contributors:* MK, RK NM: data collection; RK, SP, MK, BR: data analysis; SP, NM, MK, BR: manuscript preparation; RK, SP, BR: manuscript drafting All authors contributed to manuscript revision and its final approval.

*Funding:* None; *Competing interest:* None stated.

**WHAT THIS STUDY ADDS?**

- Utilizing specialized pediatric transport teams results in fewer transport-related adverse events in critically-ill children.

**REFERENCES**

1. Chance GW, Matthew JD, Gash J, Williams G, Cunningham K. Neonatal transport: A controlled study of skilled assistance. Mortality and morbidity of neonates less than 1.5 kg birth weight. *J Pediatr.* 1978;93:662-6.
2. Edge WE, Kanter RK, Weigle CG, Walsh RF. Reduction of morbidity in interhospital transport by specialized pediatric staff. *Crit Care Med.* 1994;22:1186-91.
3. Macnab AJ. Optimal escort for interhospital transport of pediatric emergencies. *J Trauma.* 1991;31:205-9.
4. Vos GD, Nissen AC, Nieman FH, Meurs MM, van Waardenburg DA, Ramsay G, *et al.* Comparison of interhospital pediatric intensive care transport accompanied by a referring specialist or a specialist retrieval team. *Intensive Care Med.* 2004;30:302-8.
5. Hatherill Maggie Z, Reynolds L'Argent A. Transport of critically ill children in a resource-limited setting. *Intensive Care Med.* 2003;29:1547-54.
6. Khilnani P. Available from: [http://www.pediatriconcall.com/fordocor/Conference\\_abstracts/report.aspx?reportid=450](http://www.pediatriconcall.com/fordocor/Conference_abstracts/report.aspx?reportid=450). Accessed January 08, 2015.
7. Sankar J, Singh A, Narsaria P, Dev N, Singh P, Dubey N. Prehospital transport practices prevalent among patients presenting to the pediatric emergency of a tertiary care hospital. *Indian J Crit Care Med.* 2015;19:474-8.
8. Kleinman ME, Chameides L, Schexnayder SM, Samson RA, Hazinski MF, Atkins DL, *et al.* Part 14: Pediatric Advanced Life Support 2010 American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care. *Circulation.* 2010;122:S876- S908.
9. Kumar PP, Kumar CD, Venkatlakshmi A. Long distance neonatal transport—the need of the hour. *Indian Pediatr.* 2008;45:920-2.
10. Narang M, Kaushik JS, Sharma AK, Faridi M. Predictors of mortality among the neonates transported to referral centre in Delhi, India. *Indian J Public Health.* 2013;57:100-4
11. Orr RA, Felmet KA, Han Y, McCloskey KA, Dragotta MA, Bills DM, *et al.* Pediatric specialized transport teams are associated with improved outcomes. *Pediatrics.* 2009; 124:40-8.
12. Ramnarayan P, Thiru K, Parslow RC, Harrison DA, Draper ES, Rowan KM. Effect of specialist retrieval teams on outcomes in children admitted to paediatric intensive care units in England and Wales: A retrospective cohort study. *Lancet.* 2010;376:698-704.
13. Kumar PP, Kumar CD, Shaik FA, Ghanta SB, Venkatalakshmi A. Prolonged neonatal interhospital transport on road: Relevance for developing countries. *Indian J Pediatr.* 2010;77:151-4.