

**WEB TABLE I** CLINICAL AND BIOCHEMICAL CHARACTERISTICS OF CHILDREN WITH STEROID SENSITIVE NEPHROTIC SYNDROME VS CHILDREN WITH STEROID RESISTANT NEPHROTIC SYNDROME

<i>Characteristic</i>	<i>Steroid Sensitive Nephrotic Syndrome (n=70)</i>	<i>Steroid Resistant Nephrotic Syndrome (n=30)</i>	<i>P value</i>
Age at enrolment (y)*	8 (4.2, 9.4)	7.3 (4.4, 9.3)	0.97
Age at onset of nephrotic syndrome (y)*	4.6 (1.9, 7.5)	4.6 (2, 8.4)	0.91
Duration of nephrotic syndrome (y)*	2.1 (1.2, 2.9)	2.6 (1.7, 3.6)	0.18
Male sex	44 (62)	16 (53)	0.38
Weight (kg)*	23.4 (16.7,28)	26.9 (17.5,47.9)	0.39
Height (cm) <sup>#</sup>	117.73 (20.3)	118.6 (24.7)	0.56
Body surface area (m <sup>2</sup> ) <sup>#</sup>	0.87 (0.3)	0.96 (0.32)	0.44
Serum creatinine (mg/dL) <sup>#</sup>	0.6 (0.2)	0.7 (0.3)	0.99
Serum albumin (g/dL) <sup>#</sup>	3 (0.3)	2.9 (0.4)	0.007
Serum cholesterol (mg/dL)	187 (19.3)	188.5 (29)	0.95
Spot urine protein creatinine ratio >2g/g	5 (7)	9 (30)	0.01
Spot urine protein creatinine ratio (g/g)*	0.13 (0.06,0.17)	0.16 (0.12,0.37)	0.02

*Value in n (%), \*median (IQR) or <sup>#</sup>mean (SD).*