

The isolates reported in our study were part of an on going epidemic in East Delhi. Results of phage typing of representative chloramphenicol resistant *S. typhi* strains isolated in the above study are awaited. These results are expected to throw more light on the epidemiology of these isolates.

* **V.G. Ramachandran,**
Lecturer,
Department of Microbiology
UCMS and GTB Hospital,
Delhi 110 095.

REFERENCES

1. Koul PB, Murali MV, Sharma PP, Ghai OP, Ramachandran VG, Talwar V. Multi drug resistant *Salmonella typhi* infection: Clinical Profile and therapy. *Indian Pediatr* 1991, 28: 357-361.
2. Stokes EJ. *Clinical Bacteriology*, 4th edn. London, Edwin Arnold, 1975, pp 216-222.

Reply

We appreciate the interest of Drs. Subramanyam and Lakshminarayana in our article. We agree with their point of view that disc diffusion method is influenced by many factors and ideal methods should be used.

In our study antibiotic sensitivity was done by Bauer's method(1). However, we feel the results can not be unreliable because (a) Control strains are used in this method, though on a separate plate, (b) The method is being used for over past many years, more so at many centres in the country, so emergence of sudden change in drug sensitivity pattern can not be doubted, (c) The reports of multiple drug resistant

typhoid fever from different parts of the country prove further the changing pattern of drug sensitivity in typhoid.

Harmesh Singh,
N. Raizada,
Department of Pediatrics,
Dayanand Medical College
and Hospital,
Ludhiana

REFERENCE

1. Bauer AW, Kirby WMM, Sherris TC, Trunk M. Antibiotic Susceptibility testing by a standardized single disc method. *Am J Clin Pathol* 1966, 45: 493-496.

Coin Gift : A Potentially Fatal Custom

In India, it is a common custom for a family guest to give money or coins to children of all ages in the house when he departs. This practice may be harmful to a young child as was demonstrated in this case.

A 5-month-old boy, 6 kg, presented with recurrent cough and occasional vomiting for 1 month. The child had been accepting breast feeds properly, without dysphagia or respiratory distress. Systemic examination was normal. The child has been treated symptomatically by various practitioners outside without any improvement.

In view of the lack of response of the symptoms, a chest roentgenogram was advised which revealed the presence of a coin shaped homogenous opacity in the superior mediastinum (Fig). On the lateral