

WEB TABLE I OUTCOMES AMONG CHILDREN WITH ACUTE DIARRHEA AFTER TREATMENT WITH RACECADOTRIL IN THE HOSPITAL TRIAL (N=124)

Variable	Rotavirus positive(n=58, 46.7%)			Rotavirus negative(n=66, 53.2%)		
	Racecadotril (n=31)	Placebo (n=27)	P value	Racecadotril (n=30)	Placebo (n=36)	P value
Median (IQR) duration of diarrhea (h)	26.9 (17.1,54.9)	30.2 (19, 47.1)	0.79	19.6 (14, 43.7)	23.2 (16.5,40.5)	0.45
Total volume of stool g/kg body weight: Median (IQR)	75.8 (24.6,200.8)	91.6 (9.5,194.7)	0.67	40.5 (14,118.8)	49.4 (14.6,118.8)	0.81
Average vol. of stool g/kg body weight (h): Median (IQR)	2.9 (1.7,4.7)	2.4 (0.7,4.2)	0.26	3 (1.,-4.4)	2 (0.5,3.5)	0.26
Average total fluid intake (ORS+IV) in mL g/kg body weight-h: Median (IQR)	3.6 (1.7,5.3)	3.4 (2.2,4.7)	0.84	3.6 (2.4,5.2)	2.4 (1.5,5.4)	0.30
Presence of vomiting (overall)	17 (54.8)	14 (51.9)	0.82	10 (33.3)	10 (37)	0.62
Presence of vomiting beyond 12 h; No. (%)	9 (29)	4 (14.8)	0.19	5 (16.7)	4 (11.1)	0.51
Presence of vomiting beyond 24 h; No. (%)	3 (9.7)	1 (3.7)	0.61	1 (3.3)	4 (11.1)	0.36

WEB TABLE II OUTCOMES AMONG CHILDREN WITH ACUTE DIARRHEA AFTER TREATMENT WITH RACECADOTRIL IN THE COMMUNITY TRIAL (N=196)

Variable	Rotavirus positive (n=29)			Rotavirus negative (n=167)		
	Racecadotril(n=18)	Placebo(n=11)	P value	Racecadotril (n=81)	Placebo (n= 86)	P value
Median (IQR) duration of diarrhea, d	3 (2,3)	3 (2,5)	0.48	2 (2,4)	2 (2,4)	0.83
Presence of vomiting (overall)	1 (5.6)	7 (63.6)	0.001	21 (26)	24 (28)	0.77
Presence of fever (overall)	6 (33.3)	2 (18.2)	0.67	17 (21)	16 (18.6)	0.69
Day care visit for oral rehydration; No. (%)	12 (66.7)	9 (81.8)	1.00	54 (66.7)	49 (57)	0.14