

outpatient care is time tested, however, automated dispensary systems, or pharmacy robots would reduce patient contact at the pharmacy [7]. Establishment of a telemedicine system; however, requires a robust information technology infrastructure, training of healthcare staff, receptionists, attenders, cleaning staff and security personnel; and introduction of the modifications to integrate hospital workflow. This will entail expenditure and increased cost to the patient. Development of telehealth services have in most parts of the world been hampered by the lack of insurance reimbursement for such services.

Hospitals could also consider the possibility of introducing self-administered nasal swabs for older children and adolescents; this has comparable efficacy to staff-administered swabs [8].

For children with special needs, disruptions in the schedule can be chaotic. They could be accessed through online platforms to develop home-based care programs. Teletherapy also allows the provider to document the care, train parents to maintain a regular schedule, capture sessions on video, chart progress and amend the care plan as needed.

Parents unexposed to this modality of care may be unaware of the usefulness of this form of consultation. They might feel it is impersonal and may not be satisfied with the experience. Patient acceptance can only be achieved with introduction of the often overlooked, but extremely essential operational requirement of patient education and public awareness. With these processes in place and an efficient and coordinated implementation, parents and their children would find contentment with teleconsultations like with other virtual care experiences [9].

Funding: None; *Competing interests:* None stated.

Published online: April, 2020; *PII:* S097475591600164

JEESON C UNNI

Aster Medcity, Kochi, Kerala, India.

jeeson1955@gmail.com

Do Not Neglect the Children: Considerations for COVID-19 Pandemic

The crisis of coronavirus disease (COVID-19) pandemic is the greatest challenge we have faced since World War II

REFERENCES

- Balasubramanian S, Rao NM, Goenka A, Roderick M, Ramanan AV. Coronavirus disease (covid-19) in children – what we know so far and what we do not. *Indian Pediatr.* 2020 Apr 9. Available from: https://www.indianpediatrics.net/COVID29_03.2020/SA-00159.pdf. Accessed April 24, 2020.
- Promptchara E, Ketloy C, Palaga T. Immune responses in COVID-19 and potential vaccines: Lessons learned from SARS and MERS epidemic. *Asian Pac J Allergy Immunol.* 2020;38:1-9.
- World Health Organization. Bacille Calmette-Guérin (BCG) vaccination and COVID-19. Available from: [who.int/newsroom/commentaries/detail/bacille-calmette-gu%C3%A9rin-\(bcg\)-vaccinatio-and-covid-19](http://who.int/newsroom/commentaries/detail/bacille-calmette-gu%C3%A9rin-(bcg)-vaccinatio-and-covid-19). Accessed April 24, 2020.
- Kirkpatrick BD, Colgate ER, Mychaleckyj JC, Haque R, Dickson DA, *et al.* Designed to explore complex biologic problems the “Performance of Rotavirus and Oral Polio Vaccines in Developing Countries” (PROVIDE) Study: Description of methods of an interventional study. *Am J Trop Med Hyg.* 2015;92:744-51.
- Lurie N, Carr BG. The role of telehealth in the medical response to disasters. *JAMA Intern Med.* 2018;178:745-74.
- Daniel H, Sulmasy LS. Policy Recommendations to Guide the Use of Telemedicine in Primary Care Settings: An American College of Physicians Position Paper. *Ann Intern Med.* 2015;163:787-9.
- Rodriguez-Gonzalez CG, Herranz-Alonso A, Escudero-Vilaplana V, Ais-Larigoitia MA, Iglesias-Peinado I, Sanjurjo-Saez M. Robotic dispensing improves patient safety, inventory management, and staff satisfaction in an outpatient hospital pharmacy. *Eval Clin Pract.* 2019;25:28-35.
- Akmatov MK, Gatzemeier A, Schughart, K, Pessler F. Equivalence of self- and staff collected nasal swabs for the detection of viral respiratory pathogens. *PLoS One.* 2012;7:e48508.
- Martinez KA, Rood M, Jhangiani N, Kou L, Rose S, Boissy A, *et al.* Patterns of use and correlates of patient satisfaction with a large nationwide direct to consumer telemedicine service. *J Gen Intern Med.* 2018;33:1768-73.

[1]. Although children are less commonly infected, and severity of disease in children is usually mild, no one is completely immune to this disease [2]. Most of the reported cases of COVID-19 infection in children show a complete recovery in one to two weeks; however, those with serious underlying medical conditions are at a higher risk [3].

The reason behind the lower prevalence of COVID-19

among children compared to adults is still unclear. Latest studies suggest that less exposure, less vigorous immune response, and incomplete functionality of Angiotensin-Converting Enzyme II (ACE2) in children – as the potential receptor for SARS-CoV-2 – are responsible for the significantly fewer severe cases in children [4]. Generally, children with mild symptoms and no risk factors should be managed at home [5]. Obstructed or absent breathing, central cyanosis, and cold skin are emergency signs that need immediate airway management and oxygen therapy to the target SpO₂ ≥ 94% [6]. The children with moderate and severe symptoms, along with the children with mild symptoms but with underlying risk factors, are recommended to seek hospital admission, and emergency treatment should be started according to the disease severity [5,7].

Due to a possible transmission from asymptomatic patients, children should avoid playgrounds and use a face mask in public settings. Current evidence suggests that SARS-CoV-2 is not transmittable by breast milk [8]. However, in the case of suspected or confirmed COVID-19 positive mother, feeding the expressed breast milk by a healthy caregiver is more favorable than direct breastfeeding [9]. The mother should use a mask and maintain strict hand hygiene to prevent transmission during the breast milk expressing by using alcohol-based hand rubs or soap and water. Proper disinfection should also be applied to the equipment by a healthy person.

Lockdown, along with school and playground closures, and more time spent in front of screens like video games and television, can affect children's mental health, and the lack of physical activity could result in overweight. To overcome this issue, children should be encouraged to achieve the 60-minute daily goal of physical activity, recommended by WHO, by sticking to in-door exercises [10]. Also, the quality and quantity of food should be considered seriously by the families, as the children eat unhealthier foods and the underprivileged children are undernourished due to the discontinuation of mid-day meal facility. Therefore, close supervision on physical health, nutrition, and mental health should be performed. Compared with the other groups of society, children are more vulnerable and are in need of great support [11]. It is our duty to provide support to children and families to assist them in overcoming the current situation.

Funding: None; *Competing Interests:* None stated.

Published online: April 26, 2020; *PII:* S097475591600165

AMIRREZA NASERI¹ AND MOHAMMAD-SALAR HOSSEINI^{2*}

¹*Aging Research Institute and* ²*Research Center for Evidence-Based Medicine,*

Tabriz University of Medical Sciences, Tabriz, Iran.

**hosseini.msalar@gmail.com*

REFERENCES

1. United Nations Development Program. COVID-19 pandemic: Humanity needs leadership and solidarity to defeat the coronavirus 2020. Available from: <https://www.undp.org/content/undp/en/home/coronavirus.html>. Accessed April 13, 2020.
2. Balasubramanian S, Rao NM, Goenka A, Roderick M, Ramanan AV. Coronavirus disease (covid-19) in children – what we know so far and what we do not. *Indian Pediatr.* 2020 Apr 9. Available from: <https://www.indianpediatrics.net/CONVID29.03.2020/SA-00159.pdf>. Accessed April 24, 2020.
3. Wei M, Yuan J, Liu Y, Fu T, Yu X, Zhang Z-J. Novel coronavirus infection in hospitalized infants under 1 year of age in China. *JAMA.* 2020. Available from: <https://jamanetwork.com/journals/jama/article-abstract/2761659>. Accessed April 13, 2020.
4. Dong Y, Mo X, Hu Y, Qi X, Jiang F, Jiang Z, et al. Epidemiological characteristics of 2143 pediatric patients with 2019 coronavirus disease in China. *Pediatrics.* 2020. Available from: <https://pediatrics.aappublications.org/content/pediatrics/early/2020/03/16/peds.2020-0702.full.pdf>. Accessed April 13, 2020.
5. World Health Organization (WHO). Home care for patients with COVID-19 presenting with mild symptoms and management of their contacts. Available from: [https://www.who.int/publications-detail/home-care-for-patients-with-suspected-novel-coronavirus-\(ncov\)-infection-presenting-with-mild-symptoms-and-management-of-contacts](https://www.who.int/publications-detail/home-care-for-patients-with-suspected-novel-coronavirus-(ncov)-infection-presenting-with-mild-symptoms-and-management-of-contacts). Accessed April 21, 2020.
6. World Health Organization (WHO). Clinical management of severe acute respiratory infection when COVID-19 is suspected. Available from: [https://www.who.int/publications-detail/clinical-management-of-severe-acute-respiratory-infection-when-novel-coronavirus-\(ncov\)-infection-is-suspected](https://www.who.int/publications-detail/clinical-management-of-severe-acute-respiratory-infection-when-novel-coronavirus-(ncov)-infection-is-suspected). Accessed April 21, 2020.
7. Meena J, Kumar J. Fecal shedding of SARS CoV-2: Implications for disease spread and quarantine. *Indian Pediatr.* 2020. Available from: <https://www.indianpediatrics.net/CONVID29.03.2020/CORR-00160.pdf>. Accessed April 24, 2020.
8. Chen H, Guo J, Wang C, Luo F, Yu X, Zhang W, et al. Clinical characteristics and intrauterine vertical transmission potential of COVID-19 infection in nine pregnant women: A retrospective review of medical records. *Lancet.* 2020;395:809-15.
9. Centers for Disease Control and Prevention (CDC). Considerations for inpatient obstetric healthcare settings 2020. Available from: <https://www.cdc.gov/coronavirus/2019-ncov/hcp/inpatient-obstetric-healthcare-guidance.html>. Accessed April 21, 2020.
10. World Health Organization (WHO). Global strategy on diet, physical activity and health: Activity and young people. Available from: https://www.who.int/dietphysicalactivity/factsheet_young_people/en/. Accessed April 21, 2020.
11. World Health Organization (WHO). Environmental health in emergencies: Vulnerable groups. Available from: https://www.who.int/environmental_health_emergencies/vulnerable_groups/en/. Accessed April 21, 2020.