

Impact of Water, Sanitation and Hygiene Interventions on Growth, Non-diarrheal Morbidity and Mortality in Children Residing in Low- and Middle-income Countries: A Systematic Review

TARUN GERA¹, DHEERAJ SHAH² AND HARSHPAL SINGH SACHDEV³

From ¹Department of Pediatrics, SLJain Hospital; ²Department of Pediatrics, University College of Medical Sciences (University of Delhi) & GTB Hospital; and ³Department of Pediatrics and Clinical Epidemiology, Sitaram Bhartia Institute of Science and Research; New Delhi, India.

Correspondence to: Dr Harshpal Singh Sachdev, Senior Consultant, Department of Pediatrics and Clinical Epidemiology, Sitaram Bhartia Institute of Science and Research, New Delhi, India. hpssachdev@gmail.com

Received: September 23, 2017; Initial review: November 11, 2017; Accepted: February 03, 2018.

Objective: To evaluate the impact of water, sanitation and hygiene (WASH) interventions in children (age <18 y) on growth, non-diarrheal morbidity and mortality in children.

Design: Systematic review of randomized controlled trials, non-randomized controlled trials and controlled before-after studies.

Setting: Low- and middle-income countries.

Participants: 41 trials with WASH intervention, incorporating data on 113055 children.

Intervention: Hygiene promotion and education (15 trials), water intervention (10 trials), sanitation improvement (7 trials), all three components of WASH (4 trials), combined water and sanitation (1 trial), and sanitation and hygiene (1 trial).

Outcome Measures: (i) Anthropometry: weight, height, weight-for-height, mid-arm circumference; (ii) Prevalence of malnutrition; (iii) Non-diarrheal morbidity; and (iv) mortality.

Results: There may be little or no effect of hygiene intervention on most anthropometric parameters (low- to very-low quality evidence). Hygiene intervention reduced the risk of developing Acute respiratory infections by 24% (RR 0.76; 95% CI 0.59, 0.98; moderate quality evidence), cough by 10% (RR 0.90; 95% CI 0.83, 0.97; moderate quality evidence), laboratory-confirmed influenza by 50% (RR 0.5; 95% CI 0.41, 0.62; very low quality evidence), fever by 13% (RR 0.87; 95% CI 0.74, 1.02; moderate quality evidence), and conjunctivitis by 51% (RR 0.49; 95% CI 0.45, 0.55; low quality evidence). There was low quality evidence to suggest no impact of hygiene intervention on mortality (RR 0.65; 95% CI 0.25, 1.7). Improvement in water supply and quality was associated with slightly higher weight-for-age Z-score (MD 0.03; 95% CI 0, 0.06; low quality evidence), but no significant impact on other anthropometric parameters or infectious morbidity (low to very low quality evidence). There was very low quality evidence to suggest reduction in mortality (RR 0.45; 95% CI 0.25, 0.81). Improvement in sanitation had a variable effect on the anthropometry and infectious morbidity. Combined water, sanitation and hygiene intervention improved height-for-age Z scores (MD 0.22; 95% CI 0.12, 0.32) and decreased the risk of stunting by 13% (RR 0.87; 95% CI 0.81, 0.94) (very low quality of evidence). There was no evidence of significant effect of combined WASH interventions on non-diarrheal morbidity (fever, respiratory infections, intestinal helminth infection and school absenteeism) (low- to very-low quality of evidence). Any WASH intervention (considered together) resulted in lower risk of underweight (RR 0.81; 95% CI 0.69, 0.96), stunting (RR 0.77; 95% CI 0.68, 0.86) and wasting (RR 0.12, 0.85) (low- to very-low quality of evidence).

Conclusion: Available evidence suggests that there may be little or no effect of WASH interventions on the anthropometric indices in children from low- and middle-income countries. There is low- to very-low quality of evidence to suggest decrease in prevalence of wasting, stunting and underweight. WASH interventions (especially hygiene intervention) were associated with lower risk of non-diarrheal morbidity (very low to moderate quality evidence). There was very low quality evidence to suggest some decrease to no change in mortality. These potential health benefits lend support to the ongoing efforts for provision of safe and adequate water supply, sanitation and hygiene.

Keywords: *Growth, Morbidity, Mortality, Respiratory infections, WASH interventions.*

Systematic Review Registration: PROSPERO/ CRD42017063779 (<http://www.crd.york.ac.uk/PROSPERO>)

Published online: November 11, 2017. PII:S097475591600119

The role of water supply and sanitation in controlling enteric infections, malnutrition, as well as their contribution to poverty alleviation is gaining global importance, and coverage targets for both were included in the Millennium Development Goals. Access to potable water supply and

Accompanying Editorial: Pages 377-78

proper sanitation facilities still eludes a large part of the global population, particularly in the low- and middle-income (LMIC) countries. Improvements in these aspects, also referred to as WASH (WAter supply,

Sanitation and Hygiene) interventions, are generally classified into four categories: (i) provision of an improved source of water and/or improved distribution, such as piped water or standpipes, provided either at public (source) or household (point-of-use) levels; (ii) sanitation ('hardware') interventions that provide improved means of excreta disposal; (iii) hygiene interventions that focus on health and hygiene education; and (iv) promotion of specific health behaviors like hand-washing [1].

The vast majority of research data, including systematic reviews, have focused on the impact of WASH interventions on diarrhea [2-4]. The link between WASH interventions and improvement in diarrheal infections has not translated into a demonstrable consistent improvement in other health parameters like child growth in various trials. Nutritional status of children is probably the best indicator of the health of a population, and more objective than historical recalls of diarrhea [5]. There is a paucity of systematic reviews evaluating the effect of WASH interventions on other health indicators, like malnutrition [6], mortality, and non-diarrheal morbidity; and additional trials have also now become available. We conducted this systematic review to evaluate the impact of WASH interventions on growth, non-diarrheal morbidity and mortality in children.

METHODS

Type of Studies

Individual- or cluster-randomized trials, and non-randomized and controlled before-after studies (CBA) from LMIC (individuals, families or communities) reporting outcomes in children (age <18 y) were eligible for inclusion in this review. Non-randomized trials were considered eligible for inclusion only if they had a concurrent comparison group (no WASH intervention) and adjustment for baseline characteristics and confounders. CBA studies were considered eligible for inclusion if allocation to the different comparison groups were not made by the investigators, and outcomes of interest were measured in both intervention and control groups before the WASH intervention was introduced, and again after a reasonable period of the intervention. We included non-randomized cluster trials, and CBA studies only with at least two intervention sites and two control sites.

Type of Intervention

We included studies that compared the provision of an improved source of water and/or improved distribution – such as piped water or standpipes, provided either at public (source) or household (point-of-use) levels; sanitation ('hardware') interventions that provide

improved means of excreta disposal; hygiene interventions that focused on health and hygiene education and promotion of specific health behaviors like hand-washing; and various combinations of the above listed interventions by local government, research institutions, or other non-governmental organizations – with no intervention.

Outcomes

The outcomes evaluated were: (i) anthropometry: weight, height and weight-for-height (WFH), mid-arm circumference; (ii) prevalence of malnutrition [stunting (author defined), wasting (author defined), low weight-for-age or underweight (author defined) or low BMI (author defined)]; (iii) non-diarrheal morbidity (helminth infestation, dracunculiasis, respiratory infections and others); and (iv) mortality.

Search Methods

We searched (August 2016) the following electronic databases: Medline, Web of Science, The Cochrane Controlled Trials Register, EMBASE, LILACS, Popline, and Graysource. Reference lists of all included papers and relevant reviews were scanned to identify citations that could have been missed in the primary search. We contacted authors of other relevant reviews in the field, relevant agencies and networks for the identification of ongoing or unpublished studies. The search results from the various databases and other sources were merged using reference management software (Endnote) to remove duplicate records. The title and abstract of the studies identified in the computerized search were scanned in duplicate to exclude references that were obviously irrelevant. In order to determine eligibility for inclusion of the remaining articles, their full texts were reviewed, and multiple reports of the same study were linked together. Two authors independently screened and assessed the eligibility of the studies, extracted relevant data and assessed the risk of bias for all included studies. Any dispute regarding these criteria was resolved among the investigators by mutual consultation.

Data Management

We evaluated the risk of bias for each trial using the criteria outlined in the Cochrane Handbook for Systematic Reviews of Interventions [7]. Plots of 'Risk of bias' assessments were created in Review Manager (RevMan) [8].

Risk ratio (RR) estimates with 95% confidence intervals (CI) were used for binary outcomes; for continuous outcomes, mean differences (MD) were used. In order to maximize the data input for the pooled outcome

measures, we utilized post-intervention values (means and standard deviations (SDs)) in preference to the changes from baseline [7]. In factorial trials and in multi-arm designs yielding two or more intervention groups (*e.g.*, improved water supply and promotion of hand-washing) and a single control group, the data in the intervention groups were pooled and compared against the single control group to prevent unit-of-analysis error. For cluster-randomized trials, we used the stated cluster-adjusted RR or means and 95% CI, irrespective of the method employed for adjustment. In case of missing data, we contacted trial authors for information wherever possible; and where this could not be done, or the authors did not respond, we imputed the missing values with the help of a statistician, where feasible. In case any assumptions were made for such imputations, they were recorded, and are detailed in **Web Appendix 1**.

We intended to assess contextual and clinical heterogeneity, but this was not done because of only a few

studies available for quantitative synthesis. Statistical heterogeneity was identified and measured as recommended [7]. A *P* value of 0.05 from the Chi² test was used to determine statistical significance with regard to heterogeneity. Assessment for the reporting bias using the funnel plot was also not done in view of insufficient number of trials. Subgroup analysis and sensitivity analysis were also not performed because of only a few studies available for quantitative synthesis.

We performed statistical analysis using the Revman software [8]. In concordance with the current recommendations [7], we conducted the meta-analysis of included randomized controlled trials and observational studies separately. In view of variation in studies with respect to populations, interventions, comparators, outcome and settings, the pooled effects were computed by random effects model. If it was not possible to amalgamate the data from the included studies, we provided a narrative synthesis of the results. For each primary outcome, quality




FIG. 1 The PRISMA flow chart.

assessment of the results was carried out using the GRADE approach [9].

RESULTS

The search strategy for various databases is detailed in **Web Appendix 2**, and the results are summarized in **Fig. 1**. We screened 24258 records, of which 177 were potentially eligible. Of these 107 references were excluded and 62 publications (41 studies) were included in the final analyses [10-71]. Six studies (8 publications) were ongoing [72-79].

The 41 included studies (**Web Table I**) reported data on 113055 children. Thirty-three trials were cluster randomized controlled trials, four were CBA studies, and the remaining four were cluster non-randomized controlled trials. Twenty trials were conducted in Africa, 17 in Asia and 4 in Latin America. Twenty-three trials included infants or preschool children (age <5 years), while the remaining included older children as well. The intervention was hygiene promotion and education in 15 trials, improvement in water supply, quality and storage in 10 trials, and improvement in sanitation in 7 trials. All three components of WASH intervention (water, sanitation and hygiene) were studied in 4 trials. Water and sanitation improvement was studied in one trial, and sanitation and hygiene in one study. Three trials had multiple comparison groups and yielded different combinations of interventions for analysis.

Web Fig. 1 and **Web Fig. 2** summarize the Risk of Bias for the included studies. The risk of bias for random sequence generation was low for the 33 cluster randomized controlled trials, unclear or high for two non-randomized controlled trials, and high for the remaining six studies. The risk of bias for allocation concealment was judged to be low in six, unclear in eight and high in the remaining studies. Attrition was high or unequal in the intervention and the control groups in eleven trials (high risk of bias), unclear for two trials, and low for the remaining 28 trials. Four trials were considered to be at high risk of bias on account of baseline imbalance of clusters, whereas the risk of bias was unclear for three trials, and low for remaining 34 trials. The cluster effect was not taken into account while doing the statistical analyses in six trials, and these were considered to be at high risk of bias for unit of analysis error.

Effects of Interventions

Comparison 1: Hygiene vs. No Intervention (17 trials; 82456 participants) (Table I and Web Appendix 3A)

For anthropometry, one trial [66] enrolling 1272 participants showed no evidence of difference (very low

quality evidence) in the change in anthropometry (weight, height, Z scores) between intervention and control groups (**Table I**). Two studies [16,66] evaluated the weight-for-age after long-term follow-up. Pooled analyses (no significant heterogeneity; $I^2=0%$, $P=0.9$) showed no difference (very low quality evidence) in weight-for-age or height-for-age. One trial [16] studied the impact of hygiene on BMI Z-score on follow-up and reported no change (very low quality evidence). The impact of hygiene interventions on other outcomes are also presented in **Table I**. The number of episodes of ARI were 24% lower in the hygiene intervention group ($P=0.03$; 6 trials, moderate quality evidence) (**Fig. 2**) [26,44, 50,63,65,70]. Similar benefits were also observed for cough ($P=0.006$; low quality evidence) [63], and laboratory confirmed influenza ($P<0.001$; very low quality evidence) [70]. Meta-analyses of data from four trials [50,54,58,70] showed that hygiene intervention reduced absence from school in children by 22% ($P<0.001$; moderate quality evidence). There was no evidence of any effect of hygiene intervention on mortality in children ($P=0.38$; low quality evidence).

Comparison 2: Water (Quality and Supply Improvement) vs. No Intervention

Table II presents the results of the effect of improvement in water quality and supply on various outcomes in children. Limited data from individual studies indicated marginal improvement in anthropometry, but no evidence of any significant benefit in reduction of morbidities or school absenteeism. Five trials [25,27,29,32,53] with water intervention reported on mortality data, showing a reduction in mortality by more than 50% ($P=0.007$; very low quality of evidence) (**Fig. 3** and **Web Appendix 3B**).

Comparison 3: Improvement in Sanitation vs. No Intervention

Table III presents the effect of improvement in sanitation on various outcomes. Data from individual studies did not show any significant positive effect of sanitation-related interventions on anthropometry of children, but there was a marginal benefit in terms of reduction of prevalence of underweight, wasting and stunting [13,56]. There was no evidence of significant effect on morbidity or mortality (**Fig. 4** and **Web Appendix 3C**) [23,56,69].

Comparison 4: Combined Interventions (Web Appendix 3D to 3H)

Table IV shows the magnitude of the effect in studies where more than one WASH interventions were delivered. Data on two of the WASH interventions were available only from individual studies [16,28,33,36,37,60], which did not document any significant impact on anthropometry

TABLE I EFFECT OF HYGIENE INTERVENTIONS (VS NO INTERVENTION) ON ANTHROPOMETRY, NUTRITIONAL STATUS, AND NON-DIARRHEAL MORBIDITY AND MORTALITY

Outcome	Studies	N	Effect estimate (95% CI)
Weight (kg)	1	1272	0.20 (-0.12, 0.52)*
Weight (Follow-up)	1	1390	-0.20 (-0.53, 0.13)*
Height (mm)	1	1272	10.00 (-5.39, 25.39)*
Height (Follow-up) (mm)	1	1390	-10.00 (-24.77, 4.77)*
Weight-for-age	1	1272	0.00 (-1.26, 1.26)*
WAZ (Follow-up)	2	1691	0.00 (-0.09, 0.10)*
Height-for-age	1	1272	0.00 (-0.66, 0.66)*
HAZ (Follow-up)	2	1691	-0.00 (-0.10, 0.09)#
Weight-for-Height	1	1272	0.00 (-0.99, 0.99)*
WFH (Follow-up)	1	1390	-1.00 (-1.95, -0.05)*
BMI Z-score (Follow-up)	1	301	0.10 (-0.20, 0.40)*
Low WAZ	1	168	0.85 (0.46, 1.58)\$
ARI (episodes/person-week)	6	894427	0.76 (0.59, 0.98)\$
Cough (episodes/ person-week)	1	20980	0.90 (0.83, 0.97)\$
URI (episodes/ person-week)	2	231113	0.67 (0.35, 1.28)\$
Laboratory-confirmed influenza	1	44451	0.50 (0.41, 0.62)\$
Fever	2	25140	0.87 (0.74, 1.02)\$
Skin infection	2	214293	0.80 (0.51, 1.25)\$
Conjunctivitis (episodes/person-week)	1	533416	0.49 (0.45, 0.55)\$
Intestinal parasite infection	2	1456	0.65 (0.31, 1.37)\$
School absence (episodes/person-week)	4	587825	0.78 (0.76, 0.80)\$
School absence (mean)	1	10792	0.00 (-0.01, 0.01)*
Mortality	2	5158	0.65 (0.25, 1.70)\$

*Mean difference (95% CI); #Standardized mean difference (95% CI); \$Risk ratio (95% CI); WAZ: Weight-for-age Z score; HAZ: Height-for-age Z score; WFH: Weight-for-height; ARI: Acute respiratory infection; URI: Upper respiratory infection.




FIG. 2 Forest plot of effect of Hygiene intervention versus no Intervention on incidence of acute respiratory infections (episodes/ person-week).

or morbidity. **Table V** compares the effect of any of the WASH intervention (in comparison to no intervention) on child health outcomes. There was no evidence of any significant difference in the anthropometry (weight,

height, BMI, Z scores) between the intervention and control groups, but the prevalence of underweight (**Fig. 5**), wasting and stunting (**Fig. 6**) was significantly less in intervention group [11,13,56,60].

TABLE II EFFECT OF INTERVENTIONS FOCUSING ON IMPROVEMENT IN WATER SUPPLY OR DISTRIBUTION (VS NO INTERVENTION) ON ANTHROPOMETRY, AND NON-DIARRHEAL MORBIDITY AND MORTALITY

Outcome	Studies	Participants	Effect Estimate (95% CI)
WAZ	1	121	0.03 (0.00, 0.06)*
Cough	1	5518	0.97 (0.84, 1.12)#
Fever (episodes/person-wks)	1	5518	1.02 (0.89, 1.18)#
Ocular chlamydia	1	557	1.35 (0.87, 2.09)#
Active trachoma	1	557	1.10 (0.93, 1.29)#
School absenteeism (days absent/total child-school days)	1	91946	0.99 (0.96, 1.02)#
Mortality	5	4088	0.45 (0.25, 0.81)#
RCT	4	3739	0.45 (0.25, 0.82)#
Non RCT	1	349	0.50 (0.05, 5.43)#

*Mean difference (95% CI); #Risk ratio (95% CI); WAZ: Weight-for-age Z score; RCT: Randomized controlled trial.




FIG. 3 Forest plot of effect of Water intervention on mortality.

DISCUSSION

In this systematic review of 41 trials with WASH interventions, incorporating data on 113055 children, there was no evidence of effect of hygiene intervention on anthropometry. However, hygiene intervention reduced the risk of developing acute respiratory infections by 24%, cough by 10%, laboratory-confirmed influenza by 50%, and conjunctivitis by 51%. There was low quality evidence to suggest no impact of intervention on mortality. Improvement in water supply and quality was associated with slightly higher weight-for-age Z-score

without any evidence of impact on other anthropometric measures, non-diarrheal morbidity or school absenteeism. There was very low quality evidence to suggest about 55% reduction in mortality. Improvement in sanitation had a variable effect on the anthropometry in children; no positive effect on anthropometric measures but there was a reduction in risk of wasting, stunting and underweight. Individual studies on combination of two WASH interventions did not document any significant benefit in terms of child anthropometry or morbidity. Combined water, sanitation and hygiene intervention improved height-for-age Z-scores and decreased the risk

TABLE III EFFECT OF SANITATION INTERVENTIONS (VS NO INTERVENTION) ON ANTHROPOMETRY, NUTRITIONAL STATUS, AND NON-DIARRHEAL MORBIDITY AND MORTALITY

Outcome	Studies	Participants	Effect Estimate (95% CI)
Weight	1	4315	-0.21 (-0.42, 0.01)*
Height	1	4360	-0.63 (-1.18, -0.08)*
WAZ	3	9719	-0.01 (-0.12, 0.10)*
HAZ	3	7462	-0.02 (-0.28, 0.23)*
WHZ	1	4108	-0.01 (-0.18, 0.16)*
MUAC	1	4388	-0.02 (-0.17, 0.12)*
MUAC Z-score	1	4388	0.00 (-0.13, 0.13)*
BMI Z-score	1	4104	-0.06 (-0.23, 0.11)*
Stunting	2	2791	0.88 (0.78, 0.99)#
Cluster RCT	1	2415	0.85 (0.77, 0.95)#
CBA	1	376	1.01 (0.76, 1.34)#
Underweight	2	2708	0.86 (0.76, 0.98)#
Cluster RCT	1	2452	0.85 (0.74, 0.98)#
CBA	1	256	0.98 (0.68, 1.42)#
Wasting	1	120	0.12 (0.02, 0.85)#
RTI (number of episodes)	1	5209	1.27 (1.12, 1.45)#
RTI	1	6017	0.01 (-0.02, 0.03)*
Fever	1	6015	-0.00 (-0.03, 0.02)*
Helminth infection	3	5326	0.74 (0.41, 1.33)#
Cluster RCT	2	4985	0.98 (0.86, 1.13)#
CBA	1	341	0.40 (0.28, 0.58)#
<i>C. trachomatis</i> infection	1	1211	1.01 (0.77, 1.33)#
Clinically active trachoma	2	1390	0.94 (0.83, 1.06)#
School absence (mean)	1	12262	-0.00 (-0.01, 0.01)*
Mortality (<10 years)	3	20086	1.03 (0.77, 1.39)#

*Mean difference (95% CI); #Risk ratio (95% CI); WAZ: Weight-for-age Z score; HAZ: Height-for-age Z score; WHZ: Weight-for-height Z score; MUAC: Mid upper arm circumference; RCT: Randomized controlled trial; CBA: Controlled before-after study; RTI: Respiratory tract infection.



FIG. 4 Forest plot of effect of Sanitation versus no intervention on mortality (<10 years).

of stunting. Any WASH intervention (considered together) resulted in lower prevalence of malnutrition (underweight, stunting and wasting).

Most studies in this review involved study populations from LMIC with high prevalence of malnutrition and infectious morbidities; these settings are expected to

TABLE IV EFFECT OF COMBINED (WATER, SANITATION OR HYGIENE) INTERVENTIONS (VS NO INTERVENTION) ON ANTHROPOMETRY, NUTRITIONAL STATUS, NON-DIARRHEAL MORBIDITY AND MORTALITY

<i>Outcome</i>	<i>Studies</i>	<i>Participants</i>	<i>Effect Estimate (95% CI)</i>
<i>Sanitation and Hygiene</i>			
STH	1	727	1.14 (0.87, 1.50) [#]
School absence (mean)	2	14337	-0.01 (-0.05, 0.02) [*]
<i>Water and Hygiene</i>			
WAZ (Follow-up)	1	320	-0.14 (-0.50, 0.22) [*]
HAZ (Follow-up)	1	320	-0.13 (-0.55, 0.29) [*]
BMI Z-score (Follow-up)	1	320	-0.05 (-0.39, 0.29) [*]
<i>Water and Sanitation</i>			
Low weight-for-age	1	197	0.77 (0.50, 1.19) [#]
<i>Water, Sanitation and Hygiene</i>			
HAZ	1	1899	0.22 (0.12, 0.32) [*]
Stunting	1	1899	0.87 (0.81, 0.94) [#]
STH Prevalence	2	1291	0.88 (0.60, 1.29) [#]
Cluster RCT	1	1113	1.06 (0.83, 1.36) [#]
Cluster Non-RCT	1	178	0.73 (0.57, 0.94) [#]
School absence (mean)	1	2263	-0.02 (-0.07, 0.02) [*]

^{*}Mean difference (95% CI); [#]Risk ratio (95% CI); STH: Soil transmitted helminths; WAZ: Weight-for-age Z score; HAZ: Height-for-age Z score; BMI: Body mass index; RCT: Randomized controlled trial.

benefit from WASH interventions in case of a true effect. Although the nature of interventions under each heading varied among trials, control groups in most trials were comparable with intervention groups at baseline. Thus any observed effects in the intervention groups are more likely to be attributable to the WASH strategy than to spontaneous improvements noted over time. Evidence from these trials is largely applicable to real-life situations among populations in LMIC.

Most of the studies included in this review did not have good methodological quality on some criteria. WASH is a complex intervention, and conducting field trials to evaluate its impact is challenging. By its very nature, allocation concealment and blinding of participants and observers to the intervention are very tough to execute (although a couple of trials managed to do that). Of the included trials, most were carefully conducted cluster RCTs with low risk of recruitment bias, baseline comparability of clusters, no loss of clusters and appropriate analysis. Owing to the widely varying nature of interventions, we evaluated the impact of individual group of interventions separately. This also restricted the availability of studies available for quantitative synthesis for most of the outcomes, thus downgrading the certainty of evidence for some of them.

Dangour, *et al.* [6] assessed the effect of WASH interventions on weight-for-age, weight-for-height and height-for-age Z scores. The studies included in this review were different from ours. Few studies included in this review were excluded for various reasons from the present review. In addition, we included some additional studies. The results were however similar in both the reviews, with no to minimal effect on these indices. Cumming, *et al.* [80] reviewed the effect of WASH interventions on stunting. However, it was more of a qualitative review, which focused more on observational data, and on the data from an earlier systematic review [6]. The authors suggested that WASH interventions may be effective if introduced before the onset of growth faltering. Diarrheal morbidity and mortality and onset of stunting are more concentrated before two years, and it might be important to focus on this age group to make WASH interventions more effective. A meta-analysis of the effect of hand hygiene on infectious disease risk in the community setting reported a reduction in respiratory illness of 21% (95% CI 5% to 34%) [81]. Rabie, *et al.* [82] studied the effect of handwashing on respiratory infections. All eight eligible studies reported that handwashing lowered risks of respiratory infection, with risk reductions ranging from 6% to 44% (pooled value 24%). Though none of the studies included in the review

TABLE V EFFECT OF ANY (WATER, SANITATION OR HYGIENE) INTERVENTION (VS NO INTERVENTION) ON ANTHROPOMETRY AND NUTRITIONAL STATUS OF CHILDREN

Outcome	Studies	Participants	Effect Estimate (95% CI)
Weight (kg)	2	5587	-0.02 (-0.42, 0.38)*
Weight (Follow-up)	1	1390	-0.20 (-0.53, 0.13)*
Height (mm)	2	5632	1.79 (-6.95, 10.53)*
Height (Follow-up) (mm)	1	1390	-10.00 (-24.77, 4.77)*
WAZ/WFA	5	11112	0.01 (-0.06, 0.09)#
WAZ (Follow-up)	2	2011	-0.01 (-0.10, 0.08)#
HAZ/HFA	5	10633	0.01 (-0.11, 0.14)#
HAZ (Follow-up)	2	2011	-0.01 (-0.10, 0.07)#
WFH	2	5380	-0.00 (-0.06, 0.05)#
WFH (Follow-up)	1	1390	-1.00 (-1.95, -0.05)*
MUAC	1	4388	-0.02 (-0.17, 0.12)*
MUAC Z-score	1	4388	0.00 (-0.13, 0.13)*
BMI Z-score	1	4104	-0.06 (-0.23, 0.11)*
BMI Z-score (Follow-up)	1	320	-0.05 (-0.39, 0.29)*
Underweight/Low WAZ	4	3073	0.85 (0.76, 0.97)\$
Stunting	3	4690	0.87 (0.82, 0.93)\$
Wasting	1	120	0.12 (0.02, 0.85)\$

*Mean difference (95% CI); #Standardized mean difference (95% CI); \$Risk ratio (95% CI); WAZ: Weight-for-age Z score; HAZ: Height-for-age Z score; WFH: Weight-for-height; MUAC: Mid upper arm circumference; BMI: Body mass index.




FIG. 5 Forest plot of effect of any WASH Intervention on risk of underweight (low weight-for-age).



FIG. 6 Forest plot of effect of any WASH Intervention on risk of stunting (low height-for-age).

by Rabie, *et al.* [82] were included in the present systematic review because all of these included participants from high-income countries (Australia,

Denmark, USA), these estimates are similar to our review. Pruss, *et al.* [83] reviewed the impact of the various environmental interventions on trachoma reduction.

WHAT IS ALREADY KNOWN?

- Interventions focusing on Water, Sanitation and Hygiene (WASH) result in reduction in incidence and risk of diarrhea.

WHAT THIS REVIEW ADDS?

- WASH interventions may lead to reduction in prevalence of wasting, stunting and underweight in low- and middle-income countries.
- WASH interventions (especially hygiene intervention) probably lowers risk of non-diarrheal morbidity.

However, this again was a qualitative review with bulk of the evidence emerging from observational studies, and the conclusions cannot be compared with this review.

Evidence from this review suggests that though there is little or no effect of WASH interventions on the anthropometric indices in children from LMIC, they may result in reduction in prevalence of wasting, stunting and underweight. Moreover, WASH interventions (especially hygiene intervention) are probably associated with lower risk of non-diarrheal morbidity. There are several ongoing trials on these interventions, which may alter the conclusions and improve the quality of evidence available till date. Nevertheless, these potential health benefits lend support to the ongoing efforts for provision of safe and adequate water supply, sanitation and hygiene. Future studies from varied settings need to focus on long-term benefits and other important outcomes necessary for decision-making, including the effect on micronutrient status, equity aspects and cost effectiveness.

Contributors: TG: conceptualized the review, literature search, data analysis and manuscript writing; DS: literature search data analysis and interpretation, and manuscript writing; HPS: conceptualized the review, data analysis and its interpretation, and critical inputs into manuscript writing.

Funding: Department of Health Research, Ministry of Health & Family Welfare, Government of India.

Competing interests: None stated.

REFERENCES

1. World Health Organisation (WHO)/United Nations Children's Fund (UNICEF). Global water supply and sanitation assessment 2000 report. Geneva: Water Supply and Sanitation Collaborative Council, WHO/UNICEF, 2000.
2. Clasen TF, Bostoen K, Schmidt WP, Boisson S, Fung ICH, Jenkins MW, *et al.* Interventions to improve disposal of human excreta for preventing diarrhoea. *Cochrane Database Syst Rev.* 2010;6:CD007180.
3. Fewtrell L, Kauffman B, Kay D, Enanoria W, Haller L, Colford JM. Water, sanitation, and hygiene interventions to reduce diarrhoea in less developed countries: a systematic review and meta-analysis. *Lancet Infect Dis.* 2005;5:42-52.
4. Curtis V, Cairncross S. Effect of washing hands with soap on diarrhoea risk in the community: A systematic review. *Lancet Infect Dis.* 2003;3:275-81.
5. Schmidt WP, Cairncross S. Household water treatment in poor populations: is there enough evidence for scaling up now? *Environ Sci Technol.* 2009;43:986-92.
6. Dangour AD, Watson L, Cumming O, Boisson S, Che Y, Velleman Y, *et al.* Interventions to improve water quality and supply, sanitation and hygiene practices, and their effects on the nutritional status of children. *Cochrane Database Syst Rev.* 2013;8:CD009382.
7. Higgins JPT, Green S. *Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0* (updated March 2011). The Cochrane Collaboration, 2011.
8. Cochrane Community. Editorial and Publishing Policy Resource: Review Manager (RevMan). Available from: <http://community.cochrane.org/editorial-and-publishing-policy-resource/information-technology/review-manager-revman>. Accessed September 19, 2017.
9. Gradepro. GRADE's software for Summary of Findings Tables, Health Technology Assessment and Guidelines. Available from: <https://gradepro.org/>. Accessed September 19, 2017.
10. Abdou A, Munoz BE, Nassirou B, Kadri B, Moussa F, Baarè I, *et al.* How much is not enough? A community randomized trial of a Water and Health Education programme for Trachoma and Ocular C. trachomatis infection in Niger. *Trop Med Int Health.* 2010;15:98-104.
11. Ahmed NU, Zeitlin MF, Beiser AS, Super CM, Gershoff SN, Ahmed MA. Assessment of the impact of a hygiene intervention on environmental sanitation, childhood diarrhoea, and the growth of children in rural Bangladesh. *Food Nutr Bull.* 1994;15:40-52.
12. Ahmed NU, Zeitlin MF, Beiser AS, Super CM, Gershoff SN. A longitudinal study of the impact of behavioural change intervention on cleanliness, diarrhoeal morbidity and growth of children in rural Bangladesh. *Soc Sci Med.* 1993;37:159-71.
13. Belizario Jr VY, Liwanag HJC, Naig JRA, Chua PLC, Madamba MI, Dahildahil RO. Parasitological and nutritional status of school-age and preschool-age children in four villages in Southern Leyte, Philippines: Lessons for monitoring the outcome of Community-Led Total Sanitation. *Acta Tropica.* 2015;141:16-24.

14. Boisson S, Kiyombo M, Sthreshley L, Tumba S, Makambo J, Clasen T. Field assessment of a novel household-based water filtration device: A randomised, placebo-controlled trial in the democratic republic of congo. *PloS One*. 2010;5:e12613.
15. Boisson S, Stevenson M, Shapiro L, Kumar V, Singh LP, Ward D, *et al.* Effect of household-based drinking water chlorination on diarrhoea among children under five in Orissa, India: A double-blind randomised placebo-controlled trial. *PloS Med*. 2013;10:e1001497.
16. Bowen A, Ma H, Ou J, Billhimer W, Long T, Mintz E, *et al.* A cluster randomized trial evaluating the effect of a hand washing-promotion program in Chinese primary schools. *Am J Trop Med Hyg*. 2007;76:1166-73.
17. Bowen A, Agboatwalla M, Luby S, Tobery T, Ayers T, Hoekstra RM. Association between intensive hand washing promotion and child development in Karachi, Pakistan: A cluster randomized controlled trial. *Arch Pediatr Adolesc Med*. 2012;166:1037-44.
18. Luby SP, Agboatwalla M, Painter J, Altaf A, Billhimer W, Keswick B, *et al.* Combining drinking water treatment and hand washing for diarrhoea prevention: A cluster randomised controlled trial. *Trop Med Int Health*. 2006;11:479-89.
19. Caruso BA, Freeman MC, Garn JV, Driebelbis R, Saboori S, Muga R, *et al.* Assessing the impact of a school-based latrine cleaning and hand washing program on pupil absence in Nyanza Province, Kenya: A cluster-randomized trial. *Trop Med Int Health*. 2014;19:1185-97.
20. Christensen G, Dentz HN, Pickering AJ, Bourdier T, Arnold BF, Colford JM, *et al.* Pilot cluster randomized controlled trials to evaluate adoption of water, sanitation and hygiene interventions and their combination in rural Western Kenya. *Am J Trop Med Hyg*. 2015;92:437-47.
21. Boisson S, Sosai P, Ray S, Routray P, Torondel B, Schmidt W, *et al.* Promoting latrine construction and use in rural villages practicing open defecation: Process evaluation in connection with a randomised controlled trial in Orissa, India. *BMC Research Notes* 2014;7:486.
22. Clasen T, Boisson S, Routray P, Ensink J, Jenkins M, Freeman M, *et al.* Assessing the health impact of improved rural sanitation: Designing and carrying out a cluster randomized, controlled trial in Orissa, India. *Am J Trop Med Hyg*. 2012;87(Suppl 1):231.
23. Clasen T, Boisson S, Routray P, Torondel B, Bell M, Cumming O, *et al.* Effectiveness of a rural sanitation programme on diarrhoea, soil-transmitted helminth infection, and child malnutrition in Odisha, India: A cluster-randomised trial. *Lancet Glob Health* 2014;2:e645-53.
24. Clasen T, Boisson S, Routray P, Torondel B, Jenkins M, Freeman M. The effectiveness of a rural sanitation on health and Orissa, India: A cluster-randomized, controlled trial. *Am J Trop Med Hyg*. 2014;91(Suppl 1):215.
25. Conroy RM, Meegan ME, Joyce T, McGuigan K, Barnes J. Solar disinfection of water reduces diarrhoeal disease: an update. *Arch Dis Child*. 1999;81:337-8.
26. Correa JC, Pinto D, Salas LA, Camacho JC, Rondón M, Quintero J. A cluster-randomized controlled trial of hand rubs for prevention of infectious diseases among children in Colombia. *Rev Panam Salud Publica*. 2012;31:476-484.
27. Crump JA, Oteino PO, Slutsker L, Keswick BH, Rosen DH, Hoekstra RM, *et al.* Household based treatment of drinking water with flocculant-disinfectant for preventing diarrhoea in areas with turbid source water in rural western Kenya: Cluster randomised controlled trial. *BMJ*. 2005;331:478-81.
28. Dumba R, Kaddu JB, Wabwire-Mangen F. Design and implementation of participatory hygiene and sanitation transformation (PHAST) as a strategy to control soil-transmitted helminth infections in Luweero, Uganda. *African Health Sci*. 2013;13:512-7.
29. du Preez M, Conroy R, Ligondo S, Hennessy J, Elmore-Meegan M, Soita A, *et al.* Randomized intervention study of solar disinfection of drinking water in the prevention of dysentery in Kenyan children aged under 5 Years. *Environ Sci Tech*. 2011;45:9315-23.
30. Emerson PM, Lindsay SW, Alexander N, Bah M, Dibba SM, Faal HB, *et al.* Role of flies and provision of latrines in trachoma control: Cluster-randomised controlled trial. *Lancet*. 2004;363:1093-8.
31. Emerson PM, Lindsay SW, Walraven GEL, Dibba SM, Lowe KO, Bailey RL. The Flies and Eyes Project Design and methods of a cluster-randomised intervention study to confirm the importance of flies as trachoma vectors in The Gambia and to test a sustainable method of fly control using pit latrines. *Ophthalmic Epidemiology*. 2002;9:105-17.
32. Ercumen A, Naser AM, Unicomb L, Arnold BF, Colford Jr. JM, Luby SP. Effects of source- versus household contamination of tubewell water on child diarrhea in rural Bangladesh: A randomized controlled trial. *PLoS One*. 2015;10:e0121907.
33. Fenn B, Bulti AT, Nduna T, Duffield A, Watson F. An evaluation of an operations research project to reduce childhood stunting in a food-insecure area in Ethiopia. *Public Health Nutr*. 2012;15:1746-54.
34. Driebelbis R, Freeman MC, Greene LE, Saboori S, Rheingans R. Reductions in diarrhea and clinic visits for diarrhea among children under the age of five associated with a school-based water supply, sanitation and hygiene intervention in western Kenya: A cluster-randomized trial. *Am J Trop Med Hyg*. 2012;87(suppl 1):382.
35. Freeman M, Clasen T, Driebelbis R, Greene L, Saboori S, Rheingans R. The impact of improved school water, sanitation and hygiene access on pupil diarrhea: A cluster randomized trial. *Am J Trop Med Hyg*. 2012;87(Suppl 1):428.
36. Freeman MC, Clasen T, Brooker SJ, Akoko DO, Rheingans R. The impact of a school-based hygiene, water quality and sanitation intervention on soil-transmitted helminth reinfection: A cluster-randomized trial. *Am J Trop Med Hyg*. 2013;89:875-83.
37. Freeman MC, Greene LE, Driebelbis R, Saboori S, Muga R, Brumback B, Rheingans R. Assessing the impact of a school-based water treatment, hygiene and sanitation programme on pupil absence in Nyanza Province, Kenya: A cluster-randomized trial. *Trop Med Int Health*. 2012;17:380-91.

38. Gungoren B, Latipov R, Regallet G, Musabaev E. Effect of hygiene promotion on the risk of reinfection rate of intestinal parasites in children in rural Uzbekistan. *Trans Roy Soc Trop Med Hyg.* 2007;101:564-9.
39. Gyorkos TW, Maheu-Giroux M, Blouin B, Casapia M. Impact of health education on soil-transmitted helminth infections in schoolchildren of the peruvian amazon: A cluster-randomized controlled trial. *PLoS Negl Trop Dis.* 2013;7:e2397.
40. Hammer J, Spears D. Village sanitation and children's human capital: Evidence from a randomized experiment by the Maharashtra government. Washington DC: World Bank, Sustainable Development Network, Water and Sanitation Program, 2013.
41. Huda TN, Unicomb L, Johnston RB, Halder AK, Sharkar MAY, Luby SP. Interim evaluation of a large scale sanitation, hygiene and water improvement programme on childhood diarrhea and respiratory disease in rural Bangladesh. *Soc Sci Med.* 2012;75:604-11.
42. Langford R, Lunn P, Panter-Brick C. Hand-Washing, Subclinical Infections, and Growth: A Longitudinal Evaluation of an Intervention in Nepali Slums. *Am J Human Biol.* 2011;23:621-9.
43. Anonymous. Hand washing with soap reduces the risk of diarrhoea in children. *Evidence-Based Healthcare & Public Health.* 2004;8:383-4.
44. Luby SP, Agboatwalla M, Feikin DR, Painter J, Billhimer W, Altaf A, *et al.* Effect of hand washing on child health: A randomised controlled trial. *Lancet.* 2005;366:225-33.
45. Luby SP, Agboatwalla M, Painter J, Altaf A, Billhimer WL, Hoekstra RM. Effect of intensive hand washing promotion on childhood diarrhea in high-risk communities in Pakistan: A randomized controlled trial. *JAMA.* 2004;291:2547-54.
46. Nnoaham KE. A handwashing intervention in a low income community in the developing world reduced disease incidence in children. *Evidenc Based Med.* 2006;11:88.
47. Clasen TF. Efficacy of hand washing and nail clipping on parasitic infection. *J Pediatr* 2015;167:1170-1.
48. Mahmud MA, Spigt M, Bezabih AM, Pavon IL, Dinant GJ, Velasco RB. Efficacy of hand washing with soap and nail clipping on intestinal parasitic infections in school-aged children: A factorial cluster randomized controlled trial. *PLoS Med.* 2015;12:e1001837.
49. Morris J, Schneeberger C, Jaron P, Moke F, Juma J, Ochieng J, *et al.* Ceramic Water Filters and reducing the burden of diarrheal disease in infants- Western Kenya, 2013. *Am J Trop Med Hyg.* 2014;91(Suppl 1):384.
50. Nicholson JA, Naeeni M, Hoptroff M, Matheson JR, Roberts AJ, Taylor D, *et al.* An investigation of the effects of a hand washing intervention on health outcomes and school absence using a randomised trial in Indian urban communities. *Trop Med Int Health.* 2014;19:284-92.
51. Odagiri M, Schriewer A, Daniels ME, Wuertz S, Smith WA, Clasen T, *et al.* Human fecal and pathogen exposure pathways in rural Indian villages and the effect of increased latrine coverage. *Water Res.* 2016;100:232-44.
52. Patil SR, Arnold BF, Salvatore AL, Briceno B, Ganguly S, Colford JM Jr, *et al.* The Effect of India's Total sanitation campaign on defecation behaviors and child health in rural Madhya Pradesh: A cluster randomized controlled trial. *PLoS Med.* 2014;11:e1001709.
53. Peletz R, Simunyama M, Sarenje K, Baisley K, Filteau S, Kelly P, *et al.* Assessing water filtration and safe storage in households with young children of HIV-Positive mothers: A randomized, controlled trial in Zambia. *PloS One.* 2012;7:e46548.
54. Pickering AJ, Davis J, Blum AG, Saclmanini J, Oyeir B, Okoth G, *et al.* Access to waterless hand sanitizer improves student hand hygiene behaviour in primary schools in Nairobi, Kenya. *Am J Trop Med Hyg.* 2013;89:411-18.
55. Pickering AJ, Alzua ML, Djebbari H. Impact of a community-led total sanitation intervention on child health in rural Mali: evidence from a cluster-randomised controlled trial. *Am J Trop Med Hyg.* 2014;91(Suppl1):215.
56. Pickering AJ, Djebbari H, Lopez C, Coulibaly M, Alzua ML. Effect of a community-led sanitation intervention on child diarrhoea and child growth in rural Mali: a cluster-randomised controlled trial. *Lancet Glob Health* 2015;3:e701-11.
57. Quick RE, Venczel LV, Mintz ED, Soletto L, Aparicio J, Kelly P, *et al.* Diarrhoea prevention in Bolivia through point-of-use water treatment and safe storage: A promising new strategy. *Epidemiol Infect.* 1999;122:83-90.
58. Rosen L, Manor O, Engelhard D, Brudy D, Rosen B, Peleg H, *et al.* Can a handwashing intervention make a difference? Results from a randomized controlled trial in Jerusalem preschools. *Prev Med.* 2006;42:27-32.
59. Rosen L, Manor O, Engelhard D, Zucker D. Design of the jerusalem hand washing study: Meeting the challenges of a preschool-based public health intervention trial. *Clin Trials.* 2006;3:376-84.
60. Schlesinger L, Weinberger J, Figueroa G, Secure T, González N, Mönckeberg F. Environmental sanitation: a nutrition intervention. *In: Nutrition Intervention Strategies in National Development.* Underwood Academy. Academic Press Inc. 1981:241.
61. Shafique S, Jalal CS, Jolly SP, Shikder H, Sellen D, Zlotkin S. Prevention of linear growth faltering among low birth weight infants in rural Bangladesh: A community-based cluster randomized trial. *Ann Nutr Metab.* 2013;63:1127-28.
62. Shafique S, Jalal CS, Jolly SP, Shikder H, Sellen DW, Zlotkin S. Effects of water-based hand sanitizers and micronutrient powders along with nutrition and hygiene education to prevent infections and linear growth faltering among low birth weight infants in Bangladesh. *Faseb J.* 2013;27.
63. Shafique S, Sellen DW, Lou W, Jalal CS, Jolly SP, Zlotkin SH. Mineral- and vitamin-enhanced micronutrient powder reduces stunting in full-term low-birth-weight infants receiving nutrition, health, and hygiene education: a 2 3 2 factorial, cluster-randomized trial in Bangladesh. *Am J Clin Nutr.* 2016;103:1357-69.
64. Shafique S, Shikder H, Jolly, SP, Jalal CSB, Sellen DW, Zlotkin SH. Reducing infectious morbidity and

- accelerating linear growth among low birth weight infants with hand sanitizers and nutrition and hygiene education in rural Bangladesh. *Faseb J.* 2012;26.
65. Slayton RB, Murphy JL, Morris J, Faith SH, Oremo J, Odhiambo A, *et al.* A cluster randomized controlled evaluation of the health impact of a novel anti microbial hand towel on the health of children under 2 years old in rural communities in Nyanza Province, Kenya. *Am J Trop Med Hyg.* 2016;94:437-44.
 66. Stanton BF, Clemens JD, Khair T. Educational intervention for altering water-sanitation behavior to reduce childhood diarrhea in urban Bangladesh: impact on nutritional status. *Am J Clin Nutr.* 1988;48:1166-72.
 67. Stanton BF, Clemens JD. An educational intervention for altering water sanitation behaviors to reduce childhood diarrhea in urban Bangladesh. *Am J Epidemiol.* 1987;125:292-301.
 68. Gebre T, Ayele B, Zerihun B, House J, Stoller N, Zhou Z, *et al.* Latrine Promotion for Trachoma: Assessment of Mortality from a Cluster-Randomized Trial in Ethiopia. *Am J Trop Med Hyg.* 2011;85:518-23.
 69. Stoller N, Gebre T, Ayele B, Zerihun M, Assefa Y, Habte D, *et al.* Efficacy of latrine promotion on emergence of infection with ocular Chlamydia trachomatis after mass antibiotic treatment: A cluster-randomized trial. *Int Health.* 2011;3:75-84.
 70. Talaat M, Afifi S, Dueger E, El-Ashry N, Marfin A, Kandeel A, *et al.* Effects of hand hygiene campaigns on incidence of laboratory-confirmed influenza and absenteeism in schoolchildren, Cairo, Egypt. *Energ Infect Dis.* 2011;17:619-25.
 71. West S, Munoz B, Lynch M, Kayongoya A, Chilangawa Z, Mmbaga BBO, *et al.* Impact of face-washing on trachoma in Kongwa, Tanzania. *Lancet* 1995;345:155-8.
 72. Arnold BF, Null C, Luby SP, Unicomb L, Stewart CP, Dewey KG, *et al.* Cluster randomised controlled trials of individual and combined water, sanitation, hygiene and nutritional interventions in rural Bangladesh and Kenya: The WASH Benefits study design and rationale. *BMJ Open.* 2013;3:e003476.
 73. Brown J, Cumming O, Bartram J, Cairncross S, Ensink J, Holcomb D, *et al.* A controlled, before-and-after trial of an urban sanitation intervention to reduce enteric infections in children: research protocol for the Maputo Sanitation (MapSan) study, Mozambique. *BMJ Open.* 2015;5:e008215.
 74. The Sanitation Hygiene Infant Nutrition Efficacy (SHINE) Trial Team. The Sanitation Hygiene Infant Nutrition Efficacy (SHINE) Trial: Rationale, design, and methods. *Clin Infect Dis.* 2015;61:S685-702.
 75. Jung S, Doh Y, Bizuneh DB, Beyene H, Seyong J, Kwon H, *et al.* The effects of improved sanitation on diarrheal prevalence, incidence, and duration in children under five in the SNNPR State, Ethiopia: Study protocol for a randomized controlled trial. *Trials.* 2016;17:204-13.
 76. Nery SV, Campbell S, Llewellyn S, Andrews R, Gray D, Traub R, *et al.* Impact of WASH and Albendazole distribution on infection with soil transmitted helminths in Timor-Leste: initial results of a cluster randomized controlled trial. *Am J Trop Med Hyg.* 2014;91:179.
 77. Nery SV, McCarthy JS, Traub R, Andrews RM, Black J, Gray D, *et al.* A cluster-randomised controlled trial integrating a community-based water, sanitation and hygiene programme, with mass distribution of albendazole to reduce intestinal parasites in Timor-Leste: the WASH for WORMS research protocol. *BMJ Open.* 2015;5:e009293.
 78. Overgaard HJ, Alexander N, Mátiz MI, Jaramillo JF, Olano VA. Diarrhea and dengue control in rural primary schools in Colombia: Study protocol for a randomized controlled trial. *Trials.* 2012;13:182.
 79. Overgaard, Mátiz MI, Jaramillo JF, Olano VA, Vargas SV, *et al.* Integrating dengue and diarrhea control in rural schools in Colombia: a cluster randomized controlled trial. *Am J Trop Med Hyg.* 2012;87(suppl 1):83-4.
 80. Cumming O, Cairncross S. Can water, sanitation and hygiene help eliminate stunting? Current evidence and policy implications. *Maternal Child Nutr.* 2016;12(suppl 1):91-105.
 81. Aiello AE, Coulborn RM, Perez V, Larson EL. Effect of hand hygiene on infectious disease risk in the community setting: a meta-analysis. *Am J Public Health.* 2008;98:1372-81.
 82. Rabie T, Curtis V. Handwashing and risk of respiratory infections: a quantitative systematic review. *Trop Med Int Health.* 2006;11:258-67.
 83. Pruss A, Mariotti SP. Preventing trachoma through environmental sanitation: a review of the evidence base. *Bull WHO.* 2000;78:258-66.



WEB FIG. 1 Risk of bias graph: review authors' judgements about each risk of bias item presented as percentages across all included studies.

	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of participants and personnel (performance bias)	Blinding of outcome assessment (detection bias)	Incomplete outcome data (attrition bias)	Selective reporting (reporting bias)	Other bias	Recruitment Bias	Baseline Imbalance	Loss of Clusters	Unit of Analysis Error	Comparability with Individually Randomized Controlled Trial
Abdou 2010	+	+	+	+	+	+	+	+	+	+	+	?
Ahmed 1994	+	+	+	+	+	+	+	+	+	+	+	?
Belizario Jr 2015	+	+	+	+	+	+	+	+	+	+	+	?
Boisson 2010	+	+	+	+	+	+	+	+	+	+	+	?
Boisson 2013	+	+	+	+	+	+	+	+	+	+	+	?
Bowen 2007	+	+	+	+	+	+	+	+	+	+	+	?
Bowen 2012	+	+	+	+	+	+	+	+	+	+	+	?
Caruso 2014	+	+	+	+	+	+	+	+	+	+	+	?
Christensen 2015	+	+	+	+	+	+	+	+	+	+	+	?
Clasen 2014	+	?	+	?	+	+	+	+	+	+	+	?
Conroy 1999	+	+	+	+	+	+	+	+	+	+	+	?
Correa 2012	+	+	+	+	+	+	+	+	+	+	+	?
Crump 2005	+	+	+	+	+	+	+	+	+	+	+	?
Dumba 2013	+	?	+	+	+	+	+	+	+	+	+	?
du Preez 2011	+	+	+	+	+	+	+	+	+	+	+	?
Emerson 2004	+	+	+	+	+	+	+	+	+	+	+	?
Ercumen 2015	+	+	+	+	+	+	+	+	+	+	+	?
Fenn 2012	+	+	+	+	+	+	+	+	+	+	+	?
Freeman 2013a	+	+	+	+	+	+	+	+	+	+	+	?
Gungoren 2007	+	+	+	+	+	+	+	+	+	+	+	?
Gyorkos 2013	+	?	+	+	+	+	+	+	+	+	+	?
Hammer 2013	+	+	+	+	+	+	+	+	+	+	+	?
Huda 2012	+	+	+	+	+	+	+	+	+	+	+	?
Langford 2011	?	+	+	+	+	+	+	+	+	+	+	?
Luby 2005	+	+	+	+	+	+	+	+	+	+	+	?
Mahmud 2015	+	+	+	+	+	+	+	+	+	+	+	?
Morris 2014	+	?	?	?	?	?	?	?	?	?	?	?
Nicholson 2014	+	+	+	+	+	+	+	+	+	+	+	?
Patil 2014	+	+	+	+	+	+	+	+	+	+	+	?
Peletz 2012	+	?	+	+	+	+	+	+	+	+	+	?
Pickering 2013	+	+	+	+	+	+	+	+	+	+	+	?
Pickering 2015	+	+	?	?	+	+	+	+	+	+	+	?
Quick 1999	+	+	+	+	+	+	+	+	+	+	+	?
Rosen 2006	+	+	+	+	+	+	+	+	+	+	+	?
Schlesinger 1993	+	+	+	+	+	+	+	+	+	+	+	?
Shafique 2016	+	?	+	+	+	+	+	+	+	+	+	?
Slayton 2016	+	?	+	+	+	+	+	+	+	+	+	?
Stanton 1988	+	+	+	+	+	+	+	+	+	+	+	?
Stoller 2011	+	+	+	?	+	+	+	+	+	+	+	?
Talaat 2011	+	?	+	+	+	+	+	+	+	+	+	?
West 1995	+	+	+	+	+	+	+	+	+	+	+	?

WEB FIG. 2 Risk of bias summary: review authors' judgements about each risk of bias item for each included study.

WEB APPENDIX 1 ASSUMPTIONS AND CALCULATIONS FOR IMPUTATION/INTERPRETATION OF DATA FROM STUDIES INCLUDED IN THE SYSTEMATIC REVIEWSlayton 2016

1. The total number of participants is not provided in the published paper. The number is assumed from the total number of households and the median number of two children per household.
2. The number of episodes of infection are the reported episodes of illness in the 'past' 48 hours during biweekly visits to the households.
3. The denominator of person weeks calculated by dividing the number of biweekly visits by half.

Shafique 2016

1. The data from 'Hand Sanitiser Only' and 'Hand Sanitiser and Micronutrient' groups combined as intervention and Control, and 'Micronutrient Only' groups combined as control.
2. The episodes of cough were diagnosed if the child reported to have any sort of cough or difficulty breathing. An upper respiratory tract infection was diagnosed if the mother reported symptoms of a stuffy or runny nose in her child. Both the episodes are presented separately, and combined under the section 'ARI'.

Pickering 2015

1. The infection episodes are mean proportions expressed by respondents in a 2-week recall period.
2. Respiratory infections were described under three headings: congestion, cough and difficulty breathing. Here cough is included assuming it would be a common symptom for upper and lower respiratory infections.

Patil 2014

The confidence intervals for the change in means is given. The SD is calculated with the assumption that they are the same for both the groups as calculated for Clasen 2014.

Clasen 2014

1. The individual SDs for weight-for-age and length-for-age Z-scores for the control and intervention groups was not available from the intervention data. However, the effect size (difference in mean) and its 95% CI was available. The SD for the two groups was calculated from 95% CI or SE as per standard statistical recommendations.
2. For Soil Transmitted Helmentiasis, the mean prevalence of the entire population is given. The prevalence was assumed to be the same for children and numbers calculated from there.

Pickering 2013

The number of school absence was calculated from the percentages given. Only one week data was given and this is represented as such in the calculations.

Freeman 2013a

1. A total of 1113 students were assessed at follow up. The split numbers of intervention and control groups were not provided. However, the median populations at the time of second follow up were given as 302 for intervention and 275 for control group. The total population was split in the same proportion.

For intervention, $n = (302/577) * 1113 = 583$

For control, $n = 530$

2. Only the percentage prevalence of helminth infection was mentioned. The individual numbers were calculated from percentages.
3. The same procedure was followed for school absence
4. SE converted to SD. Pupil reported absence used for quantitative analyses

Peletz 2012

For weight-for-age Z scores, the mean scores at the end and P value were mentioned.

Mean intervention = -1.21; Mean control = -1.24; $P = 0.92$

n for intervention = 61; n for control = 60

From p to t: Degree of freedom $61 + 60 - 2 = 119$

$t = 2.358$ (from table)

$SE = MD/t = -1.21 + 1.24 / 2.358 = 0.03 / 2.358 = 0.0127$

$SD = 0.0127 / 0.178 = 0.071$

Bowen 2012

SD derived from 95% CI using $SD = \sqrt{(UL - LL)}$

Rosen 2006

1. Absenteeism was analyzed in terms of the percentage of days the child was absent; number of days calculated from percentage and n .
2. There were a total of 66 days of study period. Person-weeks of exposure calculated from n and this figure.

Crump 2005

This study provided the number of deaths in children less than 5 years of age but not the total number of children. Based on the inclusion criteria, the total number of children less than 2 years is provided. We have this number as the denominator while analyzing the mortality data assuming that the proportion of children between 2-5 years would be the same in the intervention and control groups.

Emerson 2004

The number of children less than 9 years and number with trachoma calculated from percentage figures.

WEB APPENDIX 2 DETAILS OF DATABASE SEARCH AND OUTPUT

<i>Database</i>	<i>Date</i>	<i>Search Strategy</i>	<i>Number of references</i>
Medline	August 26, 2016	Water (Mesh Terms) OR Drinking Water (Mesh Terms)OR Water Quality (Mesh Terms) OR Water Purification (Mesh Terms) OR Water Supply (Mesh Terms) OR Sanitation (Mesh Terms) OR Environmental Health (Mesh Terms) OR Sanitary Engineering (Mesh Terms) OR Waste Disposal (Mesh Terms) OR Refuse Disposal (Mesh Terms) OR Drainage, Sanitary (Mesh Terms) OR Waste Management (Mesh Terms) OR Toilet Facilities (Mesh Terms) OR Hygiene (Mesh Terms) OR Hygiene, hand (Mesh Terms) OR Hand disinfection (Mesh Terms) Filters: Clinical Trial	4888
Web of Science (including Biosis Previews)	August 26, 2016	TOPIC: ('Water or Drinking Water or Water Quality or Water Purification or Water Supply or Sanitation or Environmental Health or Sanitary Engineering or Waste Disposal or Refuse Disposal or Drainage, Sanitary or Waste Management or Toilet Facilities or Hygiene or Hygiene, hand or Hand disinfection) Refined by: TOPIC: (child) AND TOPIC: (Clinical Trial)	4035
Cochrane Controlled Trials Register	August 26, 2016	'Water OR Drinking Water OR Water Quality OR Water Purification OR Water Supply OR Sanitation OR Environmental Health OR Sanitary Engineering OR Waste Disposal OR Refuse Disposal OR Drainage, Sanitary OR Waste Management OR Toilet Facilities OR Hygiene OR Hygiene, hand OR Hand disinfection in Keywords in Trials'	7900
Embase	August 27, 2016	'Water or Drinking Water or Water Quality or Water Purification or Water Supply or Sanitation or Environmental Health or Sanitary Engineering or Waste Disposal or Refuse Disposal or Drainage, Sanitary or Waste Management or Toilet Facilities or Hygiene or Hygiene, hand or Hand disinfection).mp. (mp=title, abstract, heading word, drug trade name, original title, device manufacturer, drug manufacturer, device trade name, keyword) limit 1 to (human and clinical trial and child <unspecified age>)	1182
LILACS	August 28, 2016	Water OR Drinking Water OR Water Quality OR Water Purification OR Water Supply OR Sanitation OR Environmental Health OR Sanitary Engineering OR Waste Disposal OR Refuse Disposal OR Drainage, Sanitary OR Waste Management OR Toilet Facilities OR Hygiene OR Hygiene, hand OR Hand disinfection as Subject Descriptor	564
Popline	August 29, 2016	Searched under Popline Topic 'Population Health and Environment' the subtopics Sanitation and Water Quality and Hygiene. It included keywords:Sanitation or Water Supply or Hygiene or Health Education or Water Quality or Disease Prevention and Control or Delivery of Health Care or Education or Slums or Community Development or Waste Management	3608
Greysource (Open Grey)	August 29, 2016	Water OR Sanitation OR Hygiene discipline:(06E - Medicine)	2081

WEB APPENDIX 3 SUMMARY OF FINDINGS TABLES

A. Hygiene Compared to No Intervention for Children

Patient or population: Children; Settings: Low- and Middle-income Countries
 Intervention: Hygiene ; Comparison: No intervention

Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk No intervention	Corresponding risk Hygiene				
Weight (kg)		The mean weight in the intervention groups was 0.2 higher (0.12 lower to 0.52 higher)		1272 (1 study)	⊕⊕⊕⊕ very low ^{1,2}	
Weight (Follow-up) (kg)		The mean weight (follow up) in the intervention groups was 0.2 lower (0.53 lower to 0.13 higher)		1390 (1 study)	⊕⊕⊕⊕ very low ^{1,2}	
Height (mm)		The mean height in the intervention groups was 10 higher (5.39 lower to 25.39 higher)		1272 (1 study)	⊕⊕⊕⊕ very low ^{1,2,3}	
Height (Follow-up) (mm)		The mean height (follow up) (mm) in the intervention groups was 10 lower (24.77 lower to 4.77 higher)		1390 (1 study)	⊕⊕⊕⊕ very low ^{1,2}	
Weight for age		The mean weight for age in the intervention groups was 0 higher (1.26 lower to 1.26 higher)		1272 (1 study)	⊕⊕⊕⊕ very low ^{1,2}	
WAZ (Follow-up)		The mean WAZ (follow-up) in the intervention groups was 0 standard deviations higher (0.09 lower to 0.1 higher)		1691 (2 studies)	⊕⊕⊕⊕ low ^{4,5}	SMD 0 (-0.09 to 0.1)
Height for age		The mean height for age in the intervention groups was 0 higher (0.66 lower to 0.66 higher)		1272 (1 study)	⊕⊕⊕⊕ very low ^{1,2}	
HAZ (Follow-up)		The mean HAZ (follow-up) in the intervention groups was 0 standard deviations higher (0.1 lower to 0.09 higher)		1691 (2 studies)	⊕⊕⊕⊕ low ^{4,5}	SMD 0 (-0.1 to 0.09)
Weight for Height		The mean weight for height in the intervention groups was 0 higher (0.99 lower to 0.99 higher)		1272 (1 study)	⊕⊕⊕⊕ very low ^{1,2}	
WFH (Follow up)		The mean WFH (follow up) in the intervention groups was 1 lower (1.95 to 0.05 lower)		1390 (1 study)	⊕⊕⊕⊕ very low ^{1,2}	
BMI Z score (Follow up)		The mean BMI Z score (follow up) in the intervention groups was 0.1 higher (0.2 lower to 0.4 higher)		301 (1 study)	⊕⊕⊕⊕ very low ^{2,6}	
Low WAZ	Study population		RR 0.85 (0.46 to 1.58)	168 (1 study)	⊕⊕⊕⊕ very low ^{7,8}	
	211 per 1000	179 per 1000 (97 to 334)				
Moderate	Study population					
	211 per 1000	179 per 1000 (97 to 333)				
ARI (ep/person-week)	Study population		RR 0.76 (0.59 to 0.98)	894427 (6 studies)	⊕⊕⊕⊕ moderate ⁹	
	48 per 1000	36 per 1000 (28 to 47)				
	Moderate					
	64 per 1000	49 per 1000 (38 to 63)				

Cough (episodes/ person-week)	Study population	RR 0.9 (0.83 to 0.97)	20980 (1 study)	⊕⊕⊕⊖ moderate ⁸
	118 per 1000 106 per 1000 (98 to 114)			
	Moderate			
URI (episodes/ person-week)	Study population	RR 0.67 (0.35 to 1.28)	231113 (2 studies)	⊕⊕⊖⊖ low ^{8,10}
	170 per 1000 114 per 1000 (59 to 217)			
	Moderate			
Lab Confirmed Influenza	Study population	RR 0.5 (0.41 to 0.62)	44451 (1 study)	⊕⊖⊖⊖ very low ⁸
	12 per 1000 6 per 1000 (5 to 7)			
	Moderate			
Fever	Study population	RR 0.87 (0.74 to 1.02)	25140 (2 studies)	⊕⊕⊕⊖ moderate ¹¹
	66 per 1000 57 per 1000 (49 to 67)			
	Moderate			
Skin Infection	Study population	RR 0.8 (0.51 to 1.25)	214293 (2 studies)	⊕⊕⊖⊖ low ^{8,12}
	10 per 1000 8 per 1000 (5 to 13)			
	Moderate			
Conjunctivitis (ep/person-week)	Study population	RR 0.49 (0.45 to 0.55)	533416 (1 study)	⊕⊕⊖⊖ low ⁸
	4 per 1000 2 per 1000 (2 to 2)			
	Moderate			
Intestinal Parasite Infection	Study population	RR 0.65 (0.31 to 1.37)	1456 (2 studies)	⊕⊕⊕⊖ moderate ⁸
	637 per 1000 414 per 1000 (197 to 872)			
	Moderate			
School Absence (d/person-week)	Study population	RR 0.78 (0.76 to 0.8)	587825 (4 studies)	⊕⊕⊕⊖ moderate ¹³
	70 per 1000 55 per 1000 (53 to 56)			
	Moderate			
School absence (Mean)	The mean school absence (mean) in the intervention groups was 0 higher (0.01 lower to 0.01 higher)		10792 (1 study)	⊕⊖⊖⊖ very low ^{14,15}
	Study population	RR 0.65 (0.25 to 1.7)	5158 (2 studies)	⊕⊕⊖⊖ low ⁸
5 per 1000 3 per 1000 (1 to 9)				

Moderate
14 per 1000 9 per 1000 (3 to 24)

*The basis for the assumed risk (e.g. the median control group risk across studies) is provided in footnotes. The corresponding risk (and its 95% confidence interval) is based on the assumed risk in the comparison group and the relative effect of the intervention (and its 95% CI).

CI: Confidence interval; RR: Risk ratio;

GRADE Working Group grades of evidence

High quality: Further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: We are very uncertain about the estimate.

¹ Downgraded by 1 for serious risk of bias. The included trial had high risk of bias for unit of analysis error, allocation concealment and blinding.

² Downgraded by 2 for Indirectness as there is only one included study from Asia. Extrapolation to other areas and populations not possible.

³ Downgraded by 1 for imprecision. There is only included study, with few study participants and estimates have wide confidence intervals around the estimate of the effect

⁴ Downgraded by 1 for serious risk of bias. Both trials had high risk of bias for allocation concealment and blinding and one had high risk of bias for unit of analysis error

⁵ Downgraded by 1 for indirectness as both included trials are from South Asia. Extrapolation to other areas and population not possible.

⁶ Downgraded by 1 for serious risk of bias. The one included trial had high risk of bias for blinding and allocation concealment

⁷ Downgraded by 1 for serious risk of bias. The included trial had high risk of bias for recruitment, allocation concealment and blinding.

⁸ No explanation was provided

⁹ Downgraded by 1 for serious risk of bias. All trials were at high risk of bias for blinding. Two trial were considered at high risk of bias for attrition.

¹⁰ Downgraded by 1 for imprecision. There are only two studies, with estimates that have wide confidence intervals around the estimate of the effect

¹¹ Downgraded by 1 for serious risk of bias. The two included trials had high risk of bias for allocation concealment and blinding.

¹² Downgraded by 1 for serious risk of bias. One trial had high risk of bias for allocation concealment and both for blinding

¹³ Downgraded by 1 for serious risk of bias. All four trials had high risk of bias for allocation concealment and blinding. One trial had risk of bias for attrition.

¹⁴ The one included trial had high risk of bias for blinding, allocation concealment and baseline balance between clusters.

¹⁵ The only included trial is from Africa. Extrapolation to other areas and populations not possible.

B. Improvement in Water Supply and Quality Compared to No Intervention for Children

Patient or population: Children; **Settings:** Low- and Middle-income Countries
Intervention: Improvement in Water Supply and Quality; **Comparison:** No Intervention

Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	No Intervention	Improvement in Water Supply and Quality				
WAZ		The mean waz in the intervention groups was 0.03 higher (0 to 0.06 higher)		121 (1 study)	⊕⊕⊕⊖ low ^{1,2}	
Cough	Study population		RR 0.97 (0.84 to 1.12)	5518 (1 study)	⊕⊕⊕⊖ low ^{2,3}	
	122 per 1000	118 per 1000 (102 to 136)				
	Moderate					
Fever (ep/person weeks)	Study population		RR 1.02 (0.89 to 1.18)	5518 (1 study)	⊕⊖⊖⊖ very low	
	118 per 1000	120 per 1000 (105 to 139)				
	Moderate					
Ocular Chlamydia	Study population		RR 1.35 (0.87 to 2.09)	557 (1 study)	⊕⊖⊖⊖ very low ⁴	
	110 per 1000	148 per 1000 (96 to 230)				
	Moderate					
Active Trachoma	Study population		RR 1.1 (0.93 to 1.29)	557 (1 study)	⊕⊖⊖⊖ very low	
	495 per 1000	544 per 1000 (460 to 638)				
	Moderate					
School Absenteeism (days absent/total child-school days)	Study population		RR 0.99 (0.96 to 1.02)	91946 (1 study)	⊕⊕⊕⊖ low ⁵	
	144 per 1000	142 per 1000 (138 to 146)				
	Moderate					
Mortality	Study population		RR 0.45 (0.25 to 0.81)	4088 (5 studies)	⊕⊖⊖⊖ very low ^{6,7}	
	17 per 1000	8 per 1000 (4 to 14)				
	Moderate					
Mortality - RCT	Study population		RR 0.45 (0.25 to 0.82)	3739 (4 studies)	⊕⊖⊖⊖ very low ⁸	
	18 per 1000	8 per 1000 (4 to 15)				
	Moderate					

	Moderate			
	33 per 1000	15 per 1000 (8 to 27)		
Mortality - Non RCT	Study population		RR 0.5 349	⊕⊖⊖⊖
	11 per 1000	6 per 1000 (1 to 62)	(0.05 to 5.43) (1 study)	very low ⁹
	Moderate			
	12 per 1000	6 per 1000 (1 to 65)		

*The basis for the **assumed risk** (e.g. the median control group risk across studies) is provided in footnotes. The **corresponding risk** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: Confidence interval; **RR:** Risk ratio;

GRADE Working Group grades of evidence

High quality: Further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: We are very uncertain about the estimate.

¹ The one included trial was at high risk of bias for attrition, blinding, and loss of clusters and unclear risk of bias for allocation concealment and baseline balance between clusters

² There is only one included study with small sample size from Africa. Extrapolation to other areas and populations not possible

³ The one included trial had high risk of bias for attrition

⁴ The one included trial had high risk of bias for allocation concealment, blinding, and baseline imbalance between clusters

⁵ The one included trial is from a state in India, Asia. Extrapolation to other areas and populations not possible

⁶ Three trials were at high risk of bias while one had unclear risk of bias for allocation concealment. All included trials had high risk of bias for blinding. Three trials were at high risk of bias for attrition. Three trials were at high risk of bias for loss of clusters. Two had unclear risk of bias for baseline imbalance between clusters.

⁷ Studies inadequately powered to study mortality. Total number of events (deaths) very low, below the threshold rule of thumb value of 300.

⁸ Of the 4 included trials 2 were at high risk for allocation concealment, all for blinding, 2 for attrition and 2 for loss of clusters.

⁹ The one included trial had high risk of bias for allocation concealment, blinding, attrition, loss of clusters and unclear risk of bias for random sequence generation, recruitment of clusters and baseline imbalance between clusters

C. Improvement in Sanitation Compared to No intervention for Children

Patient or population: Children; **Settings:** Low- and Middle-income Countries
Intervention: Improvement in Sanitation; **Comparison:** No intervention

Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	No intervention	Improvement in Sanitation				
Weight		The mean weight in the intervention groups was 0.21 lower (0.42 lower to 0.01 higher)		4315 (1 study)	⊕⊕⊕⊕ very low ^{1,2}	
Height		The mean height in the intervention groups was 0.63 lower (1.18 to 0.08 lower)		4360 (1 study)	⊕⊕⊕⊕ very low ^{1,2}	
WAZ		The mean WAZ in the intervention groups was 0.01 lower (0.12 lower to 0.1 higher)		9719 (3 studies)	⊕⊕⊕⊕ moderate ³	
HAZ		The mean HAZ in the intervention groups was 0.02 lower (0.28 lower to 0.23 higher)		7462 (3 studies)	⊕⊕⊕⊕ moderate ³	
WHZ		The mean WHZ in the intervention groups was 0.01 lower (0.18 lower to 0.16 higher)		4108 (1 study)	⊕⊕⊕⊕ very low ^{1,2}	
MUAC		The mean MUAC in the intervention groups was 0.02 lower (0.17 lower to 0.12 higher)		4388 (1 study)	⊕⊕⊕⊕ very low ^{1,2}	
MUAC z score		The mean MUAC Z score in the intervention groups was 0 higher (0.13 lower to 0.13 higher)		4388 (1 study)	⊕⊕⊕⊕ very low ^{1,2}	
BMI Z score		The mean bmi z score in the intervention groups was 0.06 lower (0.23 lower to 0.11 higher)		4104 (1 study)	⊕⊕⊕⊕ very low ^{1,2}	
Stunting	Study population		RR 0.88 (0.78 to 0.99)	2791 (2 studies)	⊕⊕⊕⊕ moderate ⁴	
	399 per 1000	351 per 1000 (311 to 395)				
	Moderate					
	375 per 1000	330 per 1000 (292 to 371)				
Stunting - Cluster RCT	Study population		RR 0.85 (0.77 to 0.95)	2415 (1 study)	⊕⊕⊕⊕ low ⁵	
	410 per 1000	348 per 1000 (316 to 389)				
	Moderate					
	410 per 1000	349 per 1000 (316 to 389)				
Stunting - CBA	Study population		RR 1.01 (0.76 to 1.34)	376 (1 study)	⊕⊕⊕⊕ very low ^{2,4,6}	
	340 per 1000	343 per 1000 (258 to 455)				
	Moderate					
	340 per 1000	343 per 1000 (258 to 456)				
Underweight	Study population		RR 0.86 (0.76 to 0.98)	2708 (2 studies)	⊕⊕⊕⊕ moderate ⁴	
	266 per 1000	228 per 1000 (202 to 260)				

	Moderate			
	286 per 1000	246 per 1000 (217 to 280)		
Underweight - Cluster RCT	Study population		RR 0.85	2452
	260 per 1000	221 per 1000 (193 to 255)	(0.74 to 0.98)	(1 study)
	Moderate			
	260 per 1000	221 per 1000 (192 to 255)		
Underweight - CBA	Study population		RR 0.98	256
	311 per 1000	305 per 1000 (212 to 442)	(0.68 to 1.42)	(1 study)
	Moderate			
	311 per 1000	305 per 1000 (211 to 442)		
Wasting	Study population		RR 0.12	120
	212 per 1000	25 per 1000 (4 to 181)	(0.02 to 0.85)	(1 study)
	Moderate			
	213 per 1000	26 per 1000 (4 to 181)		
RTI (number of episodes)	Study population		RR 1.27	5209
	128 per 1000	163 per 1000 (143 to 186)	(1.12 to 1.45)	(1 study)
	Moderate			
	128 per 1000	163 per 1000 (143 to 186)		
RTI		The mean rti in the intervention groups was 0.01 higher (0.02 lower to 0.03 higher)		6017
				(1 study)
Fever		The mean fever in the intervention groups was 0 higher (0.03 lower to 0.02 higher)		6015
				(1 study)
Helminth Infection	Study population		RR 0.74	5326
	155 per 1000	115 per 1000 (64 to 206)	(0.41 to 1.33)	(3 studies)
	Moderate			
	164 per 1000	121 per 1000 (67 to 218)		
Helminth Infection - Cluster RCT	Study population		RR 0.98	4985
	139 per 1000	136 per 1000 (120 to 157)	(0.86 to 1.13)	(2 studies)
	Moderate			
	110 per 1000	108 per 1000 (95 to 124)		
Helminth Infection - CBA	Study population		RR 0.4	341
	420 per 1000	168 per 1000 (118 to 244)	(0.28 to 0.58)	(1 study)
	Moderate			
	420 per 1000	168 per 1000 (118 to 244)		
Chlamydia trachomatis infection	Study population		RR 1.01	1211
	146 per 1000	147 per 1000 (112 to 194)	(0.77 to 1.33)	(1 study)

	Moderate			
	146 per 1000	147 per 1000 (112 to 194)		
Clinically Active Trachoma	Study population		RR 0.94	1390
	428 per 1000	402 per 1000 (355 to 453)	(0.83 to 1.06)	(2 studies)
	Moderate			⊕⊕⊕⊕
	287 per 1000	270 per 1000 (238 to 304)		low ^{9,10,11}
School Absence (Mean)		The mean school absence (mean) in the intervention groups was 0 higher (0.01 lower to 0.01 higher)		12262
				(1 study)
				⊕⊕⊕⊕
				very low ^{5,12}
Mortality (<10 years)	Study population		RR 1.03	20086
	19 per 1000	19 per 1000 (14 to 26)	(0.77 to 1.39)	(3 studies)
	Moderate			⊕⊕⊕⊕
	7 per 1000	7 per 1000 (5 to 10)		moderate ⁹

*The basis for the **assumed risk** (e.g. the median control group risk across studies) is provided in footnotes. The **corresponding risk** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: Confidence interval; RR: Risk ratio;

GRADE Working Group grades of evidence

High quality: Further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: We are very uncertain about the estimate.

¹ The one included trial was at high risk of bias for allocation concealment and blinding

² There is only one included trial from Asia. Extrapolation to other areas and populations not possible

³ One trial was at high risk of bias for attrition and one for allocation concealment and blinding

⁴ The one included trial was at high risk for random sequence generation, allocation concealment, blinding, recruitment bias and unclear risk of bias for unit of analysis error

⁵ There is only one included trial from Africa. Extrapolation to other populations and areas not possible.

⁶ Total number of events is less than 300 (a threshold rule-of-thumb value)

⁷ One trial was at high risk of bias for allocation concealment and blinding and one for attrition.

⁸ There are widely differing estimates of the treatment effect (i.e. heterogeneity or variability in results) across studies without a plausible explanation except study design

⁹ The included trial was at high risk of bias for allocation concealment, baseline imbalance of clusters and unit of analysis error.

¹⁰ One trial was at high risk of bias for allocation concealment, blinding and attrition

¹¹ Both trials from Africa. Extrapolation to other areas and populations not possible.

¹² One trial was at high risk of bias for allocation concealment, blinding and baseline imbalance of clusters

D. Improvement in Sanitation and Hygiene Compared to No Intervention for Children

Patient or population: Children; **Settings:** Low- and Middle-income Countries
Intervention: Improvement in Sanitation and Hygiene; **Comparison:** No Intervention

Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
No Intervention Improvement in Sanitation and Hygiene						
STH	Study population		RR 1.14 (0.87 to 1.5)	727 (1 study)	⊕⊕⊕⊕ very low ^{1,2,3}	
	208 per 1000	237 per 1000 (181 to 312)				
	Moderate					
	208 per 1000	237 per 1000 (181 to 312)				
School Absence (Mean)	The mean school absence (mean) in the intervention groups was 0.01 lower (0.05 lower to 0.02 higher)			14337 (2 studies)	⊕⊕⊕⊕ moderate ⁴	

*The basis for the **assumed risk** (e.g. the median control group risk across studies) is provided in footnotes. The **corresponding risk** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: Confidence interval; RR: Risk ratio; STH: soil transmitted helminths

GRADE Working Group grades of evidence

High quality: Further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: We are very uncertain about the estimate.

¹ The one included trial was at high risk of bias for blinding, attrition and unit of analysis error

² The one included trial is from Africa. Extrapolation to other areas and populations not possible.

³ Total number of events less than rule of thumb figure of 300

⁴ Both trials at high risk of bias for allocation concealment and blinding and one for baseline imbalance of clusters

E. Improvement in Water Supply and Quality and Hygiene Compared to No Intervention for Children

Patient or population: Children; **Settings:** Low- and Middle-income Countries
Intervention: Improvement in Water Supply and Quality and Hygiene; **Comparison:** No Intervention

Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	No Intervention	Improvement in Water Supply and Quality and Hygiene				
WAZ (Follow-up)		The mean WAZ (follow up) in the intervention groups was 0.14 lower (0.5 lower to 0.22 higher)		320 (1 study)	⊕⊖⊖⊖ very low ^{1,2}	
HAZ (Follow-up)		The mean HAZ (follow up) in the intervention groups was 0.13 lower (0.55 lower to 0.29 higher)		320 (1 study)	⊕⊖⊖⊖ very low ^{1,2}	
BMI Z-score (Follow up)		The mean BMI z score (follow up) in the intervention groups was 0.05 lower (0.39 lower to 0.29 higher)		320 (1 study)	⊕⊖⊖⊖ very low ^{1,2}	

*The basis for the **assumed risk** (e.g. the median control group risk across studies) is provided in footnotes. The **corresponding risk** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: Confidence interval;

GRADE Working Group grades of evidence

High quality: Further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: We are very uncertain about the estimate.

¹ Only one included trial from urban slums in Pakistan. Extrapolation to other populations not possible.

² Total population size is less than 400 (a threshold rule-of-thumb value; using the usual I^2 and I^2 , and an effect size of 0.2 SD).

F. Improvement in Water Supply and Quality and Sanitation Compared to No Intervention for Children

Patient or population: Children; **Settings:** Low- and Middle-income Countries
Intervention: Improvement in Water Supply and Quality and Sanitation; **Comparison:** No Intervention

Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk No Intervention	Corresponding risk Improvement in Water Supply and Quality and Sanitation				
Low weight for age	Study population		RR 0.77 (0.5 to 1.19)	197 (1 study)	⊕⊖⊖⊖ very low ^{1,2,3}	
	333 per 1000	257 per 1000 (167 to 397)				
	Moderate					
	333 per 1000	256 per 1000 (166 to 396)				

*The basis for the **assumed risk** (e.g. the median control group risk across studies) is provided in footnotes. The **corresponding risk** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: Confidence interval; RR: Risk ratio;

GRADE Working Group grades of evidence

High quality: Further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: We are very uncertain about the estimate.

¹ The one included trial was at high risk of bias for random sequence generation, allocation concealment, blinding, attrition, recruitment bias, loss of clusters and unit of analysis error

² The study was conducted in an urban slum in Chile. Extrapolation to other areas and populations not possible.

³ Total number of events less than rule of thumb value of 300,

G. Improvement in Water, Sanitation and Hygiene Compared to No Intervention for Children

Patient or population: Children; Settings: Low- and Middle-income Countries; Intervention: Improvement in Water, Sanitation and Hygiene; Comparison: No Intervention						
Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	Assumed risk	Corresponding risk				
	No Intervention	Improvement in Water, Sanitation and Hygiene				
HAZ		The mean HAZ in the intervention groups was 0.22 higher (0.12 to 0.32 higher)		1899 (1 study)	⊕⊕⊕⊕ very low ^{1,2}	
Stunting	Study population		RR 0.87 (0.81 to 0.94)	1899 (1 study)	⊕⊕⊕⊕ very low ^{1,2}	
	617 per 1000	537 per 1000 (500 to 580)				
	Moderate					
STH Prevalence	Study population		RR 0.88 (0.6 to 1.29)	1291 (2 studies)	⊕⊕⊕⊕ low ^{3,4}	
	246 per 1000	217 per 1000 (148 to 318)				
	Moderate					
STH Prevalence - Cluster RCT	Study population		RR 1.06 (0.83 to 1.36)	1113 (1 study)	⊕⊕⊕⊕ very low ^{5,6}	
	179 per 1000	190 per 1000 (149 to 244)				
	Moderate					
STH Prevalence - Cluster Non RCT	Study population		RR 0.73 (0.57 to 0.94)	178 (1 study)	⊕⊕⊕⊕ very low ^{3,7}	
	675 per 1000	493 per 1000 (385 to 634)				
	Moderate					
School Absence (Mean)		The mean school absence (mean) in the intervention groups was 0.02 lower (0.07 lower to 0.02 higher)		2263 (1 study)	⊕⊕⊕⊕ low ⁵	

*The basis for the **assumed risk** (e.g. the median control group risk across studies) is provided in footnotes. The **corresponding risk** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).
CI: Confidence interval; **RR:** Risk ratio; **STH:** soil transmitted helminths

GRADE Working Group grades of evidence: **High quality:** Further research is very unlikely to change our confidence in the estimate of effect; **Moderate quality:** Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate; **Low quality:** Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate; **Very low quality:** We are very uncertain about the estimate.

¹ The included trial was at high risk of bias for randomization, allocation concealment, blinding and recruitment bias

² The trial was conducted in rural Ethiopia. Extrapolation to other areas and populations not possible.

³ The included trial was at high risk of bias for randomization, allocation concealment and blinding

⁴ 95% confidence interval (or alternative estimate of precision) around the pooled or best estimate of effect includes both 1) no effect and 2) appreciable benefit or appreciable harm with a relative risk reduction (RRR) or relative risk increase (RRI) greater than 25%.

⁵ The study took place in school children in Kenya. Extrapolation to other areas and populations not possible.

⁶ Total number of events less than rule of thumb value of 300; ⁷ The study was conducted in rural Uzbekistan. Extrapolation to other areas and populations not possible.

H. All WASH Interventions Compared to No Intervention for Growth in Children

Patient or population: Children; **Settings:** Low- and Middle-income Countries
Intervention: All WASH Interventions; **Comparison:** No Intervention

Outcomes	Illustrative comparative risks* (95% CI)		Relative effect (95% CI)	No of Participants (studies)	Quality of the evidence (GRADE)	Comments
	No Intervention	All WASH Interventions				
Weight	Assumed risk	Corresponding risk				
		The mean weight in the intervention groups was 0.02 lower (0.42 lower to 0.38 higher)		5587 (2 studies)	⊕⊕⊖⊖ low ^{1,2}	
Weight (Follow up)		The mean weight (follow up) in the intervention groups was 0.2 lower (0.53 lower to 0.13 higher)		1390 (1 study)	⊕⊖⊖⊖ very low ^{1,3}	
Height		The mean height in the intervention groups was 1.79 higher (6.95 lower to 10.53 higher)		5632 (2 studies)	⊕⊕⊖⊖ low ^{1,2}	
Height (Follow up) (mm)		The mean height (follow up) (mm) in the intervention groups was 10 lower (24.77 lower to 4.77 higher)		1390 (1 study)	⊕⊖⊖⊖ very low ^{1,3}	
WAZ/WFA		The mean WAZ/WFA in the intervention groups was 0.01 standard deviations higher (0.06 lower to 0.09 higher)		11112 (5 studies)	⊕⊕⊕⊕ moderate ⁴	SMD 0.01 (-0.06 to 0.09)
WAZ (Follow up)		The mean WAZ (follow up) in the intervention groups was 0.01 standard deviations lower (0.1 lower to 0.08 higher)		2011 (2 studies)	⊕⊕⊕⊕ moderate ¹	SMD -0.01 (-0.1 to 0.08)
HAZ/HFA		The mean HAZ/HFA in the intervention groups was 0.01 standard deviations higher (0.11 lower to 0.14 higher)		10633 (5 studies)	⊕⊕⊕⊕ moderate ⁵	SMD 0.01 (-0.11 to 0.14)
HAZ (Follow up)		The mean HAZ (follow up) in the intervention groups was 0.01 standard deviations lower (0.1 lower to 0.07 higher)		2011 (2 studies)	⊕⊕⊕⊕ moderate ¹	SMD -0.01 (-0.1 to 0.07)
WHZ/WFH		The mean WHZ/WFH in the intervention groups was 0 standard deviations higher (0.06 lower to 0.05 higher)		5380 (2 studies)	⊕⊕⊖⊖ low ^{1,2}	SMD 0 (-0.06 to 0.05)
WFH (Follow up)		The mean WFH (follow up) in the intervention groups was 1 lower (1.95 to 0.05 lower)		1390 (1 study)	⊕⊖⊖⊖ very low ^{1,3}	
MUAC		The mean MUAC in the intervention groups was 0.02 lower (0.17 lower to 0.12 higher)		4388 (1 study)	⊕⊕⊖⊖ low ⁶	
MUAC z score		The mean MUAC z score in the intervention groups was 0 higher (0.13 lower to 0.13 higher)		4388 (1 study)	⊕⊕⊖⊖ low ⁶	
BMI Z score		The mean BMI z score in the intervention groups was 0.06 lower (0.23 lower to 0.11 higher)		4104 (1 study)	⊕⊕⊖⊖ low ⁶	
BMI z score (Follow up)		The mean BMI z score (follow up) in the intervention groups was 0.05 lower (0.39 lower to 0.29 higher)		320 (1 study)	⊕⊕⊖⊖ low ⁷	
Underweight/ Low WAZ	Study population		OR 0.81 (0.69 to 0.96)	3073 (4 studies)	⊕⊕⊖⊖ low ⁸	
	266 per 1000	227 per 1000 (200 to 258)				

	Moderate			
	286 per 1000	245 per 1000 (217 to 278)		
Stunting	Study population		OR 0.77	4690
	493 per 1000	429 per 1000 (399 to 456)	(0.68 to 0.86)	(3 studies)
	Moderate			⊕⊕⊖⊖
	410 per 1000	349 per 1000 (321 to 374)		low ⁹
Wasting	Study population		RR 0.12	120
	212 per 1000	25 per 1000 (4 to 181)	(0.02 to 0.85)	(1 study)
	Moderate			⊕⊖⊖⊖
	213 per 1000	26 per 1000 (4 to 181)		very low ^{10,11,12}

*The basis for the **assumed risk** (e.g. the median control group risk across studies) is provided in footnotes. The **corresponding risk** (and its 95% confidence interval) is based on the assumed risk in the comparison group and the **relative effect** of the intervention (and its 95% CI).

CI: Confidence interval; **RR:** Risk ratio; **OR:** Odds ratio;

GRADE Working Group grades of evidence

High quality: Further research is very unlikely to change our confidence in the estimate of effect.

Moderate quality: Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate.

Low quality: Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate.

Very low quality: We are very uncertain about the estimate.

¹ One trial was at high risk of bias for unit of analysis error and both for blinding and allocation concealment

² Both trials were conducted in rural areas in South Asia. Extrapolation to other areas and populations not possible.

³ Only one trial included from rural Bangladesh. Extrapolation to other areas and populations not possible.

⁴ One trial was at high risk of bias unit of analysis error, one for loss of clusters, two for attrition, two for allocation concealment and three for blinding.

⁵ One trial was at high risk of bias for randomization and recruitment bias, one for unit of analysis error, one for attrition, three for allocation concealment and blinding.

⁶ Only one trial from rural India. Extrapolation to other areas and populations not possible.

⁷ Only one trial from urban slums in Pakistan. Extrapolation to other areas and populations not possible.

⁸ Three of the included trials were non randomized trials. One had high risk of bias for attrition, loss of clusters and unit of analysis error. Three were at high risk of bias for allocation concealment and blinding.

⁹ Two of the three included trials were at high risk for random sequence generation, recruitment bias, allocation concealment and blinding.

¹⁰ Only one trial which was nonrandomized with high risk of bias for allocation concealment, blinding, and recruitment bias

¹¹ There is only one included trial from Asia. Extrapolation to other areas and populations not possible

¹² Total number of events is less than 300 (a threshold rule-of-thumb value)

WEB TABLE I CHARACTERISTICS OF INCLUDED STUDIES

<i>Study</i>	<i>Design</i>	<i>Country</i>	<i>Continent</i>	<i>Age group</i>	<i>N</i>			<i>Intervention category</i>	<i>Intervention details</i>	<i>Outcome of interest</i>
					<i>Total</i>	<i>Intervention</i>	<i>Control</i>			
Slayton 2016	Cluster RCT	Kenya	Africa	< 2 y	738			Hygiene	Antimicrobial Hand Towel	Acute respiratory infections, self-reported fever, and skin infections in children
Shafique 2016	Cluster RCT	Bangladesh	Asia	0-12 mo	467	236	231	Hygiene	Hand Sanitiser	Stunting, Infections
Pickering 2015	Cluster RCT	Mali	Africa	< 5 y	6012	3140	2872	Sanitation	Community-led total sanitation (CLTS) uses participatory methods to eliminate the practise of open defecation in rural communities and promote building of toilets.	Respiratory tract infections, Anthropometry, Mortality
Patil 2014	Cluster RCT	India	Asia	< 5 y	5209	2600	2609	Sanitation	Subsidies for and promotion of individual household latrines that can safely confine feces (similar to Joint Monitoring Programme defined improved sanitation facilities), school sanitation and hygiene education, Anganwadi (preschool) toilets, and community sanitation complexes.	Respiratory tract infections, Anthropometry, Worm infestations
Mahmud 2015	Cluster RCT	Ethiopia	Africa	6-15 y	367	185	182	Hygiene	Handwashing, Nail clipping	Worm infestations
Ercumen 2015	Cluster RCT	Bangladesh	Asia	6 mo-5 y	1814	1209	605	Water	Safe storage and/or water treatment	Mortality
Christensen	Cluster	Kenya	Africa	4-16 mo	432	198	234	Water,	Water treatment, sanitation	Respiratory Tract

2015	RCT							Sanitation and Hygiene	improvement, health education	Infections, Fever. Growth studied (data not provided)
Belizario 2015	CBA study	Philippines	Asia	2-15 y	341	150	191	Sanitation	Community-led total sanitation (CLTS) uses participatory methods to eliminate the practise of open defecation in rural communities and promote building of toilets.	Worm infestations, Anthropometry
Nicholson 2014	Cluster RCT	India	Asia	< 5 y	1680	847	833	Hygiene	Handwashing promotion and provision of free soap	Respiratory infections, School absence
Morris 2014	Cluster RCT	Kenya	Africa	4-10 mo	240			Water	Ceramic water filters (CWFs) remove or inactivate waterborne diarrheal pathogens in drinking water through size exclusion and silver exposure.	Respiratory infections, Febrile illness
Clasen 2014	Cluster RCT	India	Asia	0-5 y	3835	1919	1916	Sanitation	Latrine promotion and construction by combining social mobilisation with a post-hoc subsidy.	Helminth infection, Weight, Height, Mortality
Caruso 2014	Cluster RCT	Kenya	Africa	School children	17564	12262	5302	1. Sanitation and Handwashing vs Control; Handwashing vs Control; Sanitation and Handwashing vs Handwashing	Sanitation: Schools received reusable hardware (buckets, brooms, hand brushes, plastic scoop), consumables (bleach, powdered soap), toilet tissue, handwashing materials, sheets for pupils to monitor latrines conditions daily and training for two	School absence

									teachers – the head teacher and health patron. Handwashing: Received powdered soap and instructions on how to make soapy water	
Pickering 2013	Cluster RCT	Nairobi	Africa	School children	1364	895	469	Hygiene	Provision of soap and water or hand sanitizers for hand hygiene	Vomiting, cough, difficulty breathing, skin rash, rhinorrhea, school absence
Hammer 2013	Cluster RCT	India	Asia	Under 5 years				Sanitation	Latrine promotion and construction by combining social mobilisation with a subsidy.	Height
Gyorkos 2013	Cluster RCT	Peru	South America	10 y	1089	518	571	Hygiene	Hygiene Education	Soil transmitted helminthiasis prevalence
Freeman 2013a	Cluster RCT	Kenya	Africa	School children	915	470	445	Water, Sanitation and Hygiene; Water and Hygiene	Hygiene promotion, water treatment technology, and sanitation infrastructure, which included commercially manufactured hand washing and drinking water storage containers and a 1-year supply of point-of-use water treatment product distributed by Population Services International with the brand name WaterGuard.	Soil transmitted helminthiasis prevalence
Dumba 2013	Cluster RCT	Uganda	Africa	Under 5 years	727	357	370	Sanitation and Hygiene	PHAST means Participatory Hygiene and Sanitation Transformation; a participatory approach	Soil transmitted helminthiasis prevalence

									that uses visual tools to stimulate the participation of people in promotion of improved hygiene and sanitation.	
Boisson 2013	Cluster RCT	India	Asia	All children	2986	1504	1482	Water	Intensive promotion campaign and free distribution of sodium dichloroisocyanurate (NaDCC) tablets	Weight-for-age Z score; school absenteeism
Peletz 2012	Cluster RCT	Zambia	Africa	<2 y	121	61	60	Water	LifeStraw Family filter and two 5-L safe storage containers.	Weight-for-age Z score, Mortality
Huda 2012	CBA study	Bangladesh	Asia	< 5 y	1000	500	500	Water, Sanitation and Hygiene	Improvements in latrine coverage and usage; access to and use of arsenic-free water; and improved hygiene practices, especially handwashing with soap.	Acute respiratory infections
Correa 2012	Cluster RCT	Colombia	South America	1-5 y	1682	749	933	Hygiene	Alcohol based hand sanitiser	Acute respiratory infections
Bowen 2012	Cluster RCT	Pakistan	Asia	< 8 y	461	301	160	1. Hygiene; 2. Water and Hygiene	10 clusters received sodium hypochlorite solution for drinking water treatment; 9 received a flocculent-disinfectant product for drinking water treatment; 10 received soap, handwashing promotion, and flocculent disinfectant for drinking water treatment; 9 received soap and handwashing promotion; and 9 served as the control group.	Weight for age Z score, Height for age Z score, Body Mass Index Z score on long term follow-up

Fenn 2012	CBA study	Ethiopia	Africa	6 mo-3 y	1899	863	1036	Water, sanitation and hygiene	Hygiene education, pit latrines, treated water	Height for age Z score, Stunting
Talaat 2011	Cluster RCT	Egypt	Africa	Median 8 y	44451	20882	23569	Hygiene	Provision of soap and water and education	Acute respiratory infection, Influenza, Conjunctivitis, School absenteeism
Stoller 2011	Cluster RCT	Ethiopia	Africa	0-9 y	1211	608	603	Sanitation	Latrine construction	Ocular chlamydia infection; Trachoma
du Preez 2011	Cluster RCT	Kenya	Africa	6 mo - 5 y	1089	555	534	Water	Solar disinfection of water	Mortality, weight for age, height for age, weight for height
Langford 2011	Cluster non-RCT	Nepal	Asia	3-12 mo	88	45	43	Hygiene	Handwashing promotion	Weight for age Z score, Height for age Z score, Weight for height Z score, cough, cold, fever
Bosisson 2010	Cluster RCT	Congo	Africa	0-15 y	190	85	105	Water	Lifestraw Family filter for water treatment	Fever, Cough
Abdou 2010	Cluster RCT	Niger	Africa	< 5 y	557	284	273	Water	Wells and Handpump	Ocular chlamydia infection; Trachoma
Gungoren 2007	Cluster non-RCT	Uzbekistan	Asia	2-14 y	178	95	83	Water, Sanitation and Hygiene	Hand washing with soap, safe disposal of feces and boiling of drinking water.	Soil transmitted helminthiasis prevalence
Bowen 2007	Cluster RCT	China	Asia	School children	3810	2545	1265	Hygiene	Handwashing promotion, soap provision	School absence, Fever, Headache, Otagia, Rhinorrhea, Conjunctivitis, Sore throat, Cough, Vomiting
Rosen 2006	Cluster RCT	Israel	Asia	Pre - school children	1029	489	540	Hygiene	Handwashign promotion, eliminating shared cups and towels	School absence
Luby 2005	Cluster RCT	Pakistan	Asia	< 15 y	4691	3163	1528	Hygiene	Handwashing promotion, soap provision	Acute respiratory infection, Pneumonia, Impetigo, Mortality

Crump 2005	Cluster RCT	Kenya	Africa	< 5 y	715	467	248	Water	Floculent disinfectant and sodium hypochlorite	Mortality
Emerson 2004	Cluster RCT	Gambia	Africa	<9 y	179	83	96	Sanitation	Latrine construction	Trachoma
Quick 1999	Cluster RCT	Bolivia	South America	< 14 y	403	199	204	Water	Point of use water chlorination and safe storage	Soil transmitted helminthiasis prevalence
Conroy 1999	Cluster non-RCT	Kenya	Africa	< 6 y	349	175	174	Water	Solar disinfection of water	Mortality
West 1995	Cluster RCT	Tanzania	Africa	1-7 y	1417	680	737	Hygiene	Facewashing	Trachoma
Ahmed 1994	Cluster non-RCT	Pakistan	Asia	0-18 mo	168	78	90	Hygiene	Hygiene education focusing on ground sanitation, personal hygiene and food hygiene	Weight for age Z score
Stanton 1988	Cluster RCT	Bangladesh	Asia	< 6 y	1390	636	754	Hygiene	Education regarding handwashing, defecation away from house and suitable disposal of waste and faeces	Weight, Height, Weight for age Z score, Height for age Z score, Weight for Height Z score
Schlesinger 1983	CBA study	Chile	South America	0-4 y	197	113	84	Water and Sanitation	Construction of a sanitary unit consisting of a kitchen, sink and lavatory with water supply	Low weight for age

CBA: Controlled before-after; RCT: Randomized controlled trial