

Measles Case Fatality Ratio in India: A Review of Community Based Studies

CR SUDFELD AND NA HALSEY*

*From the Harvard School of Public Health, Department of Epidemiology, Boston, and * Johns Hopkins Bloomberg School of Public Health, Department of International Health, Baltimore, USA.*

Correspondence to: Christopher Sudfeld, 677 Huntington Avenue, Boston, MA 02115, USA. Email: csudfeld@hsph.harvard.edu

Context: Measles remains a major cause of child mortality in India. Measles case fatality ratios (CFRs) vary substantially between countries and even within the same community over time. We present a review of Indian community-based measles CFR studies conducted from 1975 to 2008.

Evidence acquisition: PubMed, Cochrane Libraries, and all WHO databases were searched using a combination of terms. All community-based studies were abstracted into a database.

Results: We identified 25 studies with data on 27 communities. The median CFR was 1.63 per 100 cases (Q1= 0.00 and Q3= 5.06). Studies conducted after 1994 had significantly lower CFRs ($P=0.031$). Studies in rural settings had significantly higher CFRs compared to urban studies ($P=0.015$). No differences were found by study design or outbreak/endemic setting.

Conclusions: This review suggests measles CFR may be declining in India. We hypothesize that increased measles vaccination coverage is the main factor contributing to the decline. Widespread vaccination increases both the average age of infection and the proportion of total measles cases previously vaccinated. Vitamin A treatment/supplementation is also likely to have contributed. In order to further reduce measles burden in India, vaccination and vitamin A treatment/supplementation coverage should be increased and a two dose vaccine strategy should be implemented in all areas.

Keywords: *Case Fatality Rate, India, Measles, Mortality, Vaccination.*

It is estimated that 174,000 measles deaths occurred in the Southeast Asian region during 2005, with a substantial proportion of this burden in India(1). The measles case fatality ratio (CFR) can be affected by numerous host factors including: age at infection, crowding, immunosuppression, vaccination status, malnutrition, and vitamin A deficiency(2). As a result, measles CFRs vary substantially between countries and even within the same community over time.

In 1993, WHO issued a standard protocol to determine measles case fatality ratios in a community(3). Community based studies provide the best available data in the published literature on measles CFR. Studies from Indian hospitals or other health centers are likely biased, since measles cases

with complications are likely oversampled(4). Passive surveillance (case report) studies are also prone to under-reporting of measles cases and deaths.

A recent measles CFR review of community based studies was published by WHO in 2008; however, the authors did not perform an in-depth analysis of CFRs for India(5). The most recent review of measles CFR for India was published in 1994(6). Since 1994, India has increased vaccine coverage and routine vitamin A treatment was introduced. Here we present our updated systematic review of Indian CFR of measles from community based studies published 1980-2008.

METHODS

We systematically reviewed all published literature from January 1, 1980 to December 31, 2008 to

identify Indian community based measles studies with data on measles CFRs. PubMed, Cochrane Libraries, and all World Health Organization Regional Databases were searched in all languages using combinations of the terms: *India, measles, case fatality, death, and mortality*. Prospective cohort and cross-sectional studies were abstracted. Studies were included if the study participants were from a defined Indian population with data from 1975-2008. Hospital or healthcare centre based studies and passive surveillance were excluded since these populations are likely not representative of the general Indian population.

Measles disease and measles attributed deaths were classified by the authors of the included studies. Data abstracted from the studies included: a location description (State/Union territory and urban/rural study site), year, type of study, if data were collected during an outbreak, measles cases by age, and measles deaths by age. We present CFR by the specified age groups of <1 year, 1-4 years, 5-9 years and 10+ years in order to simplify study comparison. However, if study data did not allow for these groupings, we present the data as reported.

We first performed a descriptive analysis of the studies and investigated differences by study location, outbreak setting, type of study, and year. We report the median CFR by group and in parenthesis report the 1st (Q1) and 3rd (Q3) quartiles. In order to test differences between groups, we utilized the Kruskal-Wallis test, a non-parametric method for testing equality of population medians among groups(7). Study year was dichotomized by before and after 1994, as this was the year the last Indian CFR review was published. A *P* value <0.05 was considered statistically significant for all analyses. Analyses were conducted using STATA 10.0 Special Edition (STACORP, College Station, TX).

RESULTS

We identified 25 Indian community based measles CFR studies with data on 27 distinct communities from 12 States/Union territories (8-32). Two studies presented data for two distinct populations and results were entered into the database by population (26,28). Study descriptors and results are presented for the 27 community populations in **Table I**. Twenty

of the studies were cross-sectional (74.1%) and most were conducted in rural areas (81.5%). In addition, most of the studies were performed during measles outbreaks (70.0%). A total of 8247 measles cases and 218 measles attributed deaths occurred in the studies (pooled CFR=2.64%). The mean CFR was 4.27% with a range of 0.00-31.25% and the median was 1.63 (Q1=0.00 and Q3=5.06).

Next, we analyzed the data for factors associated with measles CFRs. The median CFR for prospective studies was 1.91 (Q1=0.79 and Q3=3.52), and 1.16 (Q1=0.00 and Q3=7.00) for cross-sectional studies; the difference was not significant (*P*=0.811). The CFRs for studies conducted in rural communities (median=2.79, Q1=0.20 and Q3=7.00) were significantly higher in comparison to urban studies (median=0.00, Q1=0.00 and Q3=0.00) (*P*=0.015). The median CFR for studies performed during measles outbreaks was 2.86 (Q1=0.00 and Q3=7.07), and 1.10 (Q1=0.00 and Q3=2.19) for endemic settings; the difference was not significant (*P*=0.183).

Only 6 studies with data on 7 populations separated measles CFR by age and as a result we were unable to perform a statistical analysis of trend by age; however, in **Fig.1** we present a line graph of the data (8,9,12,13,24,26). This graph suggests a decrease in measles CFR with age, but whether CFRs for <1 yrs and 1-4 yrs differ is not clear.

We also assessed changes in measles CFR over time. In **Fig.1**, we present CFRs by midpoint study year, which suggests a decline in CFR over time. We also determined that CFR for studies occurring before 1994 (median=2.71, Q1=0.20 and Q3=7.00) were significantly greater in comparison to studies conducted after 1994 (median=0.00, Q1=0.00 and Q3=1.16) (*P*=0.031).

DISCUSSION

Measles case fatality ratios are known to significantly differ between countries and vary within populations over time(5). We reviewed Indian community based measles CFR studies to investigate factors influencing CFR and changes in mortality over time. CFR data are essential for disease burden modeling and an updated review of CFR was needed.

TABLE I DESCRIPTION AND RESULTS OF INDIAN COMMUNITY BASED STUDIES CONDUCTED FROM 1975 TO 2008

Study	Location	Urban or rural	Population age	Out-break	Years	Type of study*	Total measles Cases	Total deaths	Over all CFR	<1y CFR	1-4y CFR	5-9y CFR	>10y CFR	Other age groups
Dhanoa(8)	Punjab	Rural	<2 yrs		1970s	P	82	3	3.66	2.63				1-2y 6.82
Chand(9)	Uttar Pradesh	Rural	<14 yrs		1974-86	P	411	9	2.19	7.27				1-3y 1.69 4-14y 0.83
Garai(10)	West Bengal	Rural	Children		1976-78	P	862	2	0.2					
John(11)	Tamil Nadu	Rural	<10 yrs	Yes	1977-78	P	198	14	7.07					
Cherian(12)	Tamil Nadu	Rural	<5 yrs	Yes	1979-80	C	78	8	10.26	9.09	10.45			
Vasudev(13)	Uttar Pradesh	Rural	<12 yrs		1980	P	266	9	3.38	0	6.5			
Swami(14)	Rajasthan	Urban	<15 yrs		1980-81	P	731	10	1.37					
Jajoo(15)	Maharashtra	Rural	<10 yrs	Yes	1982	P	113	0	0					
Sharma(16)	Rajasthan	Rural	<5 yrs	Yes	1982	C	88	4	4.55					
Lakhanpal(17)	Punjab	Rural	<14 yrs		1982-83	C	241	2	0.83					
Bhatia(18)	Nagaland	Rural	<12 yrs	Yes	1983	C	515	14	2.71					
Rao(19)	Karnataka	Rural	<5 yrs		1983	C	132	0	0					
Lobo(20)	Haryana	Rural	<10 yrs		1984	P	430	7	1.63					
Sharma(21)	Rajasthan	Rural	<14 yrs	Yes	1984	C	133	19	14.29					
Mangal(22)	Rajasthan	Urban	<10 yrs		1985-86	C	189	0	0					
Gupta(23)	Himachal Pradesh	Rural	<15 yrs	Yes	1986	C	217	11	5.06					
Narain(24)	Uttar Pradesh	Rural	All ages	Yes	1986	C	771	54	7	23.07	11.47	5.52	0	
Satpathy(25)	West Bengal	Rural	All ages	Yes	1986	C	581	1	0.17					
Risbud(26)	Maharashtra	Rural	<10 yrs	Yes	1991	C	48	15	31.25	37.5	33.33	0		
Risbud(26)	Maharashtra	Rural	<10 yrs	Yes	1992	C	128	20	15.63	33.33	13.04	6.67		
Thakur(27)	Chandigarh	Urban	All ages	Yes	1998-99	C	283	0	0					
John(28)	Tamil Nadu	Rural	<10 yrs	Yes	1999	C	70	2	2.86					
Ray(29)	West Bengal	Urban	<5 yrs		2000	C	290	0	0					
Sharma(30)	Chandigarh	Urban	<14 yrs	Yes	2003	C	58	0	0					
Gupta(23)	Himachal Pradesh	Rural	<15 yrs	Yes	2004	C	69	0	0					
Mishra(32)	Madhya Pradesh	Rural	Children	Yes	2004	C	1204	14	1.16					
John(28)	Tamil Nadu	Rural	<19 yrs	Yes	2006	C	59	0	0					

* P = Prospective; C = Cross-sectional.

Measles CFR in India appears to have decreased during 1975-2008. We hypothesize that increased measles vaccination coverage in India is the main factor contributing to this decline, in addition to other factors including the introduction of vitamin A in case management and increasing vitamin A supplementation coverage. However, it may not be appropriate to generalize from this review that CFR has decreased for the entire population of India, since published data are only available for select communities in 12 Indian states or Union territories. In addition, 38% of all districts in India still had measles vaccine coverage less than 50% in 2005 and these districts are not proportionally represented in this review(33).

Widespread measles vaccination increases the average age of measles infection at the population level by decreasing the force of infection(34). Data from the US Centers for Disease Control and the recent WHO measles CFR review suggest that children <5 yrs infected with measles have increased mortality in comparison to children infected at an older age(2,5). Due to the small number of Indian CFR studies reporting the age of study participants, we were not able to statistically test differences between age groups. Nevertheless, a visual analysis of the Indian data in **Fig. 2** suggests that measles CFR is decreased in children >5 years.

A single dose measles vaccine is estimated to be 85% efficacious in preventing measles disease, and

as a result a proportion of the total measles cases occurring in a community are expected to have been previously vaccinated(35). The proportion of total measles cases previously vaccinated in a community is anticipated to increase as vaccine coverage increases(36). For example, if measles vaccine coverage for a population is 50%, 13% of the total measles cases are expected to have been previously vaccinated. Whereas, if vaccination coverage is 90%, 57% of the total measles cases are expected to have been previously vaccinated. Multiple observational studies have found decreased measles mortality or measles complications in the previously vaccinated(2,37-39). There is clear evidence of partial immunity in some studies, but confounding by differential access to health care could be a factor in some studies. When measles vaccination coverage increases, the expected proportion of total cases previously vaccinated increases, and in turn, the population case fatality ratio likely decreases.

Vitamin A deficiency is a known risk factor for measles mortality(40). Since 1987, the WHO and UNICEF have recommended vitamin A treatment of children with measles(41). A meta-analysis of randomized controlled trials found 200,000 IU of vitamin A given for 2 days was associated with a 64% reduction in overall mortality(42). Nevertheless, measles case management with vitamin A may not have considerably affected Indian CFRs at the population level, since coverage of vitamin A treatment has been shown to be low in multiple




FIG.1 Measles case fatality ratio by age category.




FIG.2 Measles case fatality ratio by year of study (Studies with multiple years of follow-up are plotted at study midpoint).

communities with high levels of measles transmission. A recent observational study in Madhya Pradesh found that only 15.8% of measles cases received therapeutic doses of vitamin A and another study conducted in slum areas of Kolkata found only 8.6% were treated(19,32). Routine vitamin A supplementation is also thought to decrease measles case fatality; however, the data suggest supplementation may not be as effective in preventing measles mortality as vitamin A administration at the onset of measles(43). Coverage of vitamin A supplementation may also be low in high risk populations. In a recent study in the slums of Delhi, only 37.6% percent of children 12-23 received a vitamin A supplement(44). Vitamin A treatment and routine supplementation have likely contributed to declining CFRs in India, but due to low coverage in communities at high risk for measles disease and mortality, the impact on population CFRs may not be considerable.

The data also suggest higher CFRs in rural areas compared to urban communities. This difference may be attributable to differences in access to health care and vaccination services. No significant differences were found by study design or for studies conducted in outbreak vs. endemic settings. These findings are similar to the results of the WHO case fatality review(5).

Overall, this review suggests measles CFR may be declining in India over time. We theorize that increased measles vaccination coverage is the main contributor to the decline. The impact of increasing vaccination coverage on measles mortality is greater than that expected from prevention of measles disease alone; since at higher coverage levels, the average age of infection is older and a larger proportion of measles cases are expected to have been previously vaccinated. Vitamin A treatment and supplementation decrease an individual's risk of measles mortality, but the impact in India at the population level may be minimal due to low coverage. In order to continue to decrease measles CFR in India; measles vaccination, vitamin A treatment, and routine vitamin A supplementation coverage should be increased. In addition, the cost effective strategy of introducing supplementary immunization activities to provide children with two

doses of measles vaccine as well as increase single dose coverage could also significantly decrease mortality(45). India has greatly reduced the total number of measles cases and deaths over the past few decades, yet much more needs to be done to decrease the substantial burden of this preventable disease.

Contributors: CRS conducted literature search and data abstraction. CRS and NAH wrote the manuscript. Data abstraction and analyses were conducted while CRS was at Johns Hopkins.

Financial support: This work was supported in part by a grant to the US Fund for UNICEF from the Bill & Melinda Gates Foundation (Grant 43386).

Competing Interests: None stated.

REFERENCES

1. Wolfson LJ, Strebel PM, Gacic-Dobo M, Hoekstra EJ, McFarland JW, Hersh BS. Has the 2005 measles mortality reduction goal been achieved? A natural history modelling study. *Lancet* 2007; 369:191-200.
2. Perry RT, Halsey NA. The clinical significance of measles: a review. *J Infect Dis* 2004;189 Suppl 1:S4-16.
3. Byass P. Measles control in the 1990s: generic protocol for determining measles case fatality rates in a community, either during an epidemic or in a high endemic area. (WHO/EPI/GEN/93.3). Geneva: WHO; 1993.
4. Narain JP, Banerjee KB. Measles in India: epidemiology and control. *Indian J Pediatr* 1989; 56: 463-472.
5. Wolfson LJ, Grais RF, Luquero FJ, Birmingham ME, Strebel PM. Estimates of measles case fatality ratios: a comprehensive review of community-based studies. *Int J Epidemiol* 2009; 38: 192-205.
6. Singh J, Sharma RS, Verghese T. Measles mortality in India: a review of community based studies. *J Commun Dis* 1994; 26: 203-214.
7. Kruskal WH, Wallis WA. Use of ranks in one-criterion variance analysis. *J Amer Statist Assn* 1952; 47: 583-621.
8. Dhanoa J, Cowan B. Measles in the community- a study in non hospitalised young children in Punjab. *J Trop Pediatr* 1982; 28: 59-61.
9. Chand P, Rai RN, Chawla U, Tripathi KC, Datta

- KK. Epidemiology of measles—a thirteen years prospective study in a village. *J Commun Dis* 1989; 21: 190-199.
10. Garai R, Chakraborty AK. Measles in a rural community. *Indian J Public Health* 1981; 24: 150-153.
 11. John TJ, Joseph A, George TI, Radhakrishnan J, Singh RP, George K. Epidemiology and prevention of measles in rural south India. *Indian J Med Res* 1980; 72: 153-158.
 12. Cherian T, Joseph A, John TJ. Low antibody response in infants with measles and children with subclinical measles virus infection. *J Trop Med Hyg* 1984; 87: 27-31.
 13. Vasudev JP, Nandan D, Chandra R, Srivastava BC. Post measles complications in a rural population. *J Commun Dis* 1983; 15: 249-252.
 14. Swami SS, Chandra S, Dudani IU, Sharma R, Mathur MM. Epidemiology of measles in western Rajasthan. *J Commun Dis* 1987; 19: 370-372.
 15. Jajoo UN, Chhabra S, Gupta OP, Jain AP. Measles epidemic in rural community near Sevagram. *Indian J Public Health* 1984; 28: 204-207.
 16. Sharma RS, Kaushic VK, Johri SP, Ray SN. An epidemiological investigation of measles outbreak in Alwar-Rajasthan. *J Commun Dis* 1984; 16: 299-303.
 17. Lakhanpal U, Rathore MS. Epidemiology of measles in rural area of Punjab. *J Commun Dis* 1986; 18: 185-188.
 18. Bhatia R. Measles outbreak in village Tophema in Nagaland. *J Commun Dis* 1985; 17: 185-189.
 19. Rao RS, Kumari J, Rao TS, Narashimham VL. Measles in a rural community. *J Commun Dis* 1988; 20: 131-135.
 20. Lobo J, Reddaiah VP, Kapoor SK, Nath LM. Epidemiology of measles in a rural community. *Indian J Pediatr* 1987; 54: 261-265.
 21. Sharma RS. An epidemiological study of measles epidemic in district Bhilwara, Rajasthan. *J Commun Dis* 1988; 20: 301-311.
 22. Mangal N, Shah K, Sitaraman S. Epidemiological study of measles in urban (slum) area of Jaipur. *Indian Pediatr* 1990; 27: 1216-1217.
 23. Gupta BP, Swami HM, Bhardwaj AK, Vaidya NK, Sharma CD, Kaushal RK. An outbreak of measles in a remote tribal area of Himachal Pradesh. *Indian J Comm Health* 1989; 5: 25-28.
 24. Narain JP, Khare S, Rana SR, Banerjee KB. Epidemic measles in an isolated unvaccinated population, India. *Int J Epidemiol* 1989; 18: 952-958.
 25. Satpathy SK, Chakraborty AK. Epidemiological study of measles in Singur, West Bengal. *J Commun Dis* 1990; 22: 23-26.
 26. Risbud AR, Prasad SR, Mehendale SM, Mawar N, Shaikh N, Umrani UB, *et al.* Measles outbreak in a tribal population of Thane district, Maharashtra. *Indian Pediatr* 1994; 31: 543-551.
 27. Thakur JS, Ratho RK, Bhatia SP, Grover R, Issaivanan M, Ahmed B, *et al.* Measles outbreak in a periurban area of Chandigarh: need for improving vaccine coverage and strengthening surveillance. *Indian J Pediatr* 2002; 69: 33-37.
 28. John S, Sanghi S, Prasad S, Bose A, George K. Two doses of measles vaccine: are some states in India ready for it? *J Trop Pediatr* 2009; 55: 253-256.
 29. Ray SK, Mallik S, Munsri AK, Mitra SP, Baur B, Kumar S. Epidemiological study of measles in slum areas of Kolkata. *Indian J Pediatr* 2004; 71: 583-586.
 30. Sharma MK, Bhatia V, Swami HM. Outbreak of measles amongst vaccinated children in a slum of Chandigarh. *Indian J Med Sci* 2004; 58: 47-53.
 31. Gupta BP, Sharma S. Measles Outbreak in a rural area near Shimla. *Indian J Community Med.* 2006; 31: 106-108.
 32. Mishra A, Mishra S, Jain P, Bhadoriya RS, Mishra R, Lahariya C. Measles related complications and the role of vitamin A supplementation. *Indian J Pediatr* 2008; 75: 887-890.
 33. Department of Family Welfare, Ministry of Health and Family Welfare. Multi Year Strategic Plan 2005-2010. Universal Immunization Programme. Government of India; 2005.
 34. Grenfell BT, Anderson RM. The estimation of age-related rates of infection from case notifications and serological data. *J Hyg (Lond)* 1985; 95: 419-436.
 35. Cutts FT, Grabowsky M, Markowitz LE. The effect of dose and strain of live attenuated measles vaccines on serological responses in young infants. *Biologicals* 1995; 23: 95-106.

36. Orenstein WA, Bernier RH, Dondero TJ, Hinman AR, Marks JS, *et al.* Field evaluation of vaccine efficacy. *Bull World Health Organ* 1985; 63: 1055-1068.
 37. Aaby P, Bukh J, Lisse IM, da Silva MC. Decline in measles mortality: nutrition, age at infection, or exposure? *Br Med J (Clin Res Ed)* 1988; 296: 1225-1228.
 38. Byass P, Adedeji MD, Mongdem JG, Zwandor AC, Brew-Graves SH, Clements CJ. Assessment and possible control of endemic measles in urban Nigeria. *J Public Health Med* 1995;17: 140-145.
 39. Hull HF, Williams PJ, Oldfield F. Measles mortality and vaccine efficacy in rural West Africa. *Lancet* 1983;1: 972-75.
 40. Sommer A, West KP. *Vitamin A deficiency: health, survival, and vision.* New York: Oxford University Press; 1996.
 41. WHO. Joint WHO/UNICEF statement on vitamin A for measles. *International Nursing Review* 1988; 35: 21.
 42. D'Souza RM, D'Souza R. Vitamin A for the treatment of children with measles—a systematic review. *J Trop Pediatr* 2002; 48: 323-327.
 43. The Vitamin A and Pneumonia Working Group. Potential interventions for the prevention of childhood pneumonia in developing countries: A meta-analysis of data from field trials to assess the impact of Vitamin A supplementation on pneumonia morbidity and mortality. *Bull WHO* 1995; 73: 609-619.
 44. Sachdeva S, Datta U. Vitamin A-first dose supplement coverage evaluation amongst children aged 12-23 months residing in slums of Delhi, India. *Indian J Ophthalmol* 2009; 57: 299-303.
 45. Dabral M. Cost effectiveness of supplementary immunization for measles in India. *Indian Pediatr* 2009; 46: 957-962.
-