

Drug Induced Acute Dystonic Reaction

A 5-year-old, developmentally normal boy, presented with sudden onset involuntary, persistent, deviation of neck towards left, along with protrusion of tongue and hoarse cry for last 5 hours (**Fig. 1** and **Webvideo 1**). It was associated with neck pain and inability to close the mouth. There was history of fever for 1 day and he received four doses of oral paracetamol (15 mg/kg/dose) over a period of 24 hours. Examination showed spasmodic torticollis to left, persistent open jaw, and protruded tongue. Rest of the neurological examination was essentially normal. Acute onset drug-induced dystonic crisis was considered and child was treated with single intravenous dose of chlorpheniramine (0.2 mg/kg); prompt response was seen with subsidence of dystonic reaction within next 20 minutes. He was observed for next 24 hours, and there was no recurrence of dystonic crisis.

Acute dystonic crisis can occur with antidepressants, dopamine receptor blocking agents, antiemetics and antipsychotics. Drug-induced dystonic reaction commonly present as acute onset focal dystonia characterized by torticollis, tongue protrusion and laryngeal spasm. Paracetamol is a selective inhibitor of cyclooxygenase and in usual doses it does not cross the blood brain barrier. Higher doses may activate central serotonergic pathways resulting in central cholinergic and dopaminergic imbalance. Acute onset cervical dystonia is very unusual with therapeutic dosage of paracetamol.



FIG.1 Spasmodic torticollis to left.

Treatment include immediate withdrawal of offending agent and anticholinergic agents like chlorpheniramine, benzodiazepines or dopaminergic agonists. Acute dystonic reaction is often misdiagnosed as seizure, encephalitis or tetany. Eliciting a through history is important to avoid unnecessary investigations and treatment for this potentially reversible condition.

INDAR KUMAR SHARAWAT AND RENU SUTHAR*

Pediatric Neurology Unit,

Department of Pediatrics, PGIMER, Chandigarh, India.

**drrenusuthar@gmail.com*

NOTICE

Call for Submission of 'Clinical Videos'

Under this section, *Indian Pediatrics* publishes videos depicting an intricate technique or an interesting clinical manifestation, which are difficult to describe clearly in text or by pictures. A video file submitted for consideration for publication should be of high resolution and should be edited by the author in final publishable format. MPEG or MP4 formats are acceptable. The maximum size of file should be 20 MB. The file should not have been published elsewhere, and will be a copyright of *Indian Pediatrics*, if published. For this section, there should be a write-up of up to 250 words discussing the condition and its differential diagnoses. The write-up should also be accompanied by a thumbnail image for publication in the print version and PDF. Submit videos as separate Supplementary files with your main manuscript. A maximum of three authors (not more than two from a single department) are permissible for this section. In case the video shows a patient, he/she should not be identifiable. In case the identification is unavoidable, or even otherwise, each video must be accompanied by written permission of parent/guardian, as applicable. Authors are responsible for obtaining participant consent-to-disclose forms for any videos of identifiable participants, and should edit out any names mentioned in the recording. The consent form should indicate its purpose (publication in the journal in print and online, with the understanding that it will have public access) and the signed consent of the parent/legal guardian. The copy of the consent form must be sent as supplementary file along with the write-up, and original form should be retained by the author. A sample consent form is available at our website www.indianpediatrics.net.